

Taxonomy of the Genus *Pseudonaja* (Reptilia: Elapidae) in Australia.

by

Richard W. Wells

“Shiralee”, Major West Road, Cowra, New South Wales, Australia

The clear morphological differences that exist within the genus as previously considered strongly indicate that it is a polyphyletic assemblage. Accordingly, I have taken the step of formally proposing the fragmentation of *Pseudonaja*. In this work I have decided to restrict the genus *Pseudonaja* to the *Pseudonaja nuchalis* complex. Additionally, I herein formally resurrect from synonymy the generic name *Euprepiosoma* Fitzinger, 1860 for the *textilis* group of species, erect a new generic name (*Placidaserpens* gen. nov.) for the snakes previously regarded as *Pseudonaja guttata*, erect a new generic name (*Notopseudonaja* gen. nov.) for the group of species previously regarded as the *Pseudonaja modesta* complex, and erect a new generic name (*Dugitophis* gen. nov.) for snakes previously regarded as the *Pseudonaja affinis* complex.

Genus *Pseudonaja* Gunther, 1858 The *Pseudonaja nuchalis* Complex

It is usually reported that *Pseudonaja nuchalis* occurs across most of northern, central and western Australia, ranging from Cape York Peninsula, in the north-east, through western, southern and south-eastern Queensland, far western New South Wales, north-western Victoria, and most of South Australia, Northern Territory and Western Australia. However, this distribution pattern is now known to actually represents several different species all regarded by most authorities for convenience as the single highly variable species, '*Pseudonaja nuchalis*'. As usually defined, this actually is a highly variable and therefore confusing group of species to identify and it is not all surprising that there has been difficulty in breaking up the group. Another not insignificant reason behind this reticence to look at '*Pseudonaja nuchalis*' more closely, is that all are highly venomous, very fast and numerous fatalities have resulted from their bites. Until recent years, virtually any description of this 'species' would state or infer that the colour and pattern is subject to considerable ontogenetic, geographic and even seasonal variation, and to an extent this is still true for the *Pseudonaja nuchalis* complex. Snakes presently regarded as 'Western Brown Snakes' in most texts have a basic body colour that can range across most shades of brown, right through to black, but creamish, yellow, orange and reddish variations are known. Some specimens with the orange or brown base colour are often totally unpatterned, or just barely marked with a few black scales on the neck (often arranged in a 'W' or 'V' pattern), while others can have the head and/or neck greyish, brownish or even jet black, with an unpatterned or patterned body. Patterning can vary from merely scattered black dorsal scales, to neat reticulations of dark-edged scales, to even broad dark rings, bands or blotches. Regardless of the dorsal colour and

patterning, the ventral surface is usually light yellowish to pale orange or creamish with irregular rows of orange spots or small blotches in most areas.

The colour of the buchal cavity is purplish-black. Juveniles usually have a generalised pattern of blackish head and neck patches and a body pattern of faint reticulations, but some juveniles have similar patterns to their respective adult 'variations'. As can be seen from this range of colour and pattern combinations, the 'Western Brown Snake' has been widely recorded over Australia since its original description. Morphologically, the scalation patterns appear conservative across the different taxa, so it may not be possible to identify the different taxa alone on the basis of a scale-count. However, recent investigators have now come to the conclusion that the 'Western Brown Snake' is in fact not one highly variable species, but several different species. To field naturalists this has not been all that much of a surprise for the differences in behaviour, habitat and morphology between the various populations more than hinted that a number of different species were in fact being lumped into the name *Pseudonaja nuchalis*. This snake has now been subjected to a range of morphological, genetic and biochemical investigations by some of the world's leading scientists (see references), and it is now finally gaining wider acceptance that several different species have been unknowingly included under the name '*Pseudonaja nuchalis*' in the past. Some of these 'variations' are included in this work as different species partly as a consequence of their distinctive chromosomal arrangements following the work of Mengden (1985).

Various estimates are that '*Pseudonaja nuchalis*' may actually represent anywhere from 4 to in excess of 10 different species. In this work *Pseudonaja nuchalis* has been split into 8 different species - and all are supported by chromosomal evidence. Some (but by no means all) of the numerous 'variations' of this species often called 'Western Brown Snakes' are as follows:

Firstly, the snake that has been traditionally regarded as the 'Western Brown Snake' *Pseudonaja nuchalis* is in fact based on a specimen from near Arnhem Land. It is in actuality the most highly restricted of all the group and really would be more appropriately called the 'Northern Brown Snake' instead. Its colour pattern is one of striking wide dark bands on a yellowish-orange to orange-brown base colour.

The variation previously known as the 'southern morph' of *Pseudonaja nuchalis*, should now be called *Pseudonaja aspidorhyncha*. In this species the body colour may be any shade of uniform brown, and in most cases the body is unpatterned. Usually the head is slightly darker brown than the body, and there can be a few scattered black scales on the nape. In occasional specimens the head can be very dark brown and the body may have scattered but faint darker brown scales; in some individuals the head can be completely black.

In another 'morph' (now called *Pseudonaja kellyi*), the head and neck region are jet black (or sometimes very dark brown). The base colour of the body is yellowish-orange to orange-brown, with many of the dorsal scales marked with dark brown or black to form a transverse ziz-zag or even a 'herring-bone' pattern over the body. Juveniles have a similar colour and pattern to the adults, with the exception that the snout is paler followed by a darker interocular area and a dark brownish nape.

Another 'morph' (now called *Pseudonaja imperitor*), has the base colour of the body uniform tan-brown, and the ventral surface creamish. The snout is pale creamish-brown, followed by an interocular band of dark brown over the head. The nuchal area can have a few scattered black scales, or occasionally these can form a narrow band over the neck. Juveniles have a similar colour and pattern to the adults, with the exception that the darker interocular area and nape markings are more pronounced. There is quite pronounced seasonal colour change with this form - during the summer the entire body colour becomes much lighter brown, while during the (slightly) cooler winter (or dry season) it changes to a darker brown overall.

Another 'morph' (now called *Pseudonaja mengdeni*) has a base body colour varying from tan-brown through to pale yellow or orange.

There is a strong 'herring bone' pattern on the posterior two-thirds of the body caused through the arrangement of darker reddish-brown scales. The head and neck is usually pale brown (snout is paler) with a slightly darker interocular area.

The neck is greyish-brown to darker brown, with a dark narrow row of black or very dark brown scales often forming a sharp boundary (sometimes in a 'V' shape) immediately anterior to the nape patch. Juveniles have a similar colour and pattern to the adults, with the exception that the darker interocular area and nape markings are more pronounced.

Another very distinctive 'morph' (now called *Pseudonaja carinata*) has the head brown, with the nape paler containing a few darker scales. The base body colour is pale creamish-brown, with the posterior of the body heavily banded with up to 13 (mostly 11) broad blotches or saddle-like bands of black. The dark blotches are slightly narrower than the paler interspaces anteriorly, but are as wide as or slightly wider than the pale areas posteriorly. Within the pale interspaces there are three or four narrow (1 scale wide) dark brown bands. Juveniles have a similar colour and pattern to the adults, with the exception that the head is darker and the broad dark body bands either completely encircle the body or break around the medial area of the ventrals. The WA population of this 'morph' appears to have a greater number of bands also and this could indicate that it may be taxonomically distinct in itself.

In still another 'morph' (now called *Pseudonaja acutirostris*) the base body colour is brownish to pale orange, with the body being strongly banded in black. Sometimes the anterior of the body is unpatterned, with exception of a couple of black scales on the neck or occasionally a blackish patch on the nape. More often however, the body has a series of (up to 14) very broad black bands, each of which is usually narrower than the paler interspaces. Within the paler areas there are a few thin faint bands of reddish-brown.

Another 'morph' (now called *Pseudonaja gowi*) has an overall plain brown body colour and reduced patterning. However, with this type there is the addition of a broad black band near the neck-nuchal area, and this is occasionally accompanied by a thin secondary collar of black at the anterior end of the neck band. Additionally, there can be a loose vertebral series of black scales on the posterior of the body. The ventral area is creamish with pale orange blotching.

There are many other variations that are apparently rarely encountered, some of which may represent other undescribed species, hybrids, or merely just colour variations of some of the above. This is a group of snakes that still requires urgent study.

Mitchell's Brown Snake

***Pseudonaja acutirostris* (Mitchell, 1951)**

Previously known as the 'southern, orange with black bands morph', of *Pseudonaja nuchalis*, genetic studies by Mengden et al have proven its distinctiveness from that species, necessitating the re-instatement of the original name given it by Mitchell in 1955 - thus it should now be called *Pseudonaja acutirostris*.

Diagnosis: This is a medium to large species with a relatively slender body, and a small narrow head that is not distinct from the neck. The eye is large with a round pupil and a pale iris. As presently understood, this species has a distinctive chromosomal morphology (of the $2n=34$ karyotype). Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. The base body colour is brownish to pale orange, with the body being strongly banded in black.

Sometimes the anterior of the body is unpatterned, with exception of a couple of black scales on the neck or occasionally a blackish patch on the nape. More often however, the body has a series of (up to 14) very broad black bands, each of which is usually narrower than the paler interspaces. Within the paler areas there are a few thin faint bands of reddish-brown. It attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m.

Notes: This species has a scattered distribution over arid and semi-arid eastern and central Australia, ranging from western New South Wales, north-western Victoria, eastern and northern South Australia, and the southern part of the Northern Territory, and probably adjacent parts of Western Australia. Its principle habitat is open woodland and shrubs with scattered grass cover on open plains with stoney soils and in places low rocky hills. As in the case with all members of the *Pseudonaja nuchalis* complex, it is an oviparous species, producing up to 20 eggs in a clutch. The main diet comprises lizards but small mammals may also be taken. This snake is highly venomous and although there are no records of fatalities, urgent medical attention should be sought in the event of a bite, because this is most certainly a potentially dangerous species. Protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Additionally, it is protected under the Victorian Wildlife Act (1975) [but not listed as threatened in Schedule 2 of the Victorian Flora and Fauna Guarantee Act (1988)], the SA National Parks and Wildlife Act (1972), the Territory Parks and Wildlife Conservation Act (1998), and the WA Wildlife Conservation Act 1950 (as amended). Generally this species is regarded as being common over much of its range, but in Victoria its survival status may be classified as of 'Lower Risk - Near Threatened' because of its limited occurrence in that State. Further, it is possible that its present distribution pattern elsewhere suggests that its range has fragmented, so it may be considered as potentially vulnerable in some areas.

McCoy's Brown Snake

***Pseudonaja aspidorhyncha* (McCoy, 1879)**

Previously known as the 'southern morph' of *Pseudonaja nuchalis*, recent genetic studies have proven its distinctiveness from that species, necessitating the re-instatement of the original name given it by Frederick McCoy in 1879 - thus it should now be called *Pseudonaja aspidorhyncha*.

Diagnosis: This is a large and relatively slender snake with a small narrow head not distinct from the neck, the eye is large with a round pupil and a pale reddish iris. As presently understood, this species has a distinctive chromosomal morphology (of the $2n=34$ karyotype), and could actually represent at a minimum a polytypic species in itself. Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. The body colour may be any shade of uniform brown, and in most cases the body is unpatterned. The head is slightly darker brown than the body, and there can be a few scattered black scales on the nape. In occasional specimens the head can be very dark brown and the body may have scattered but faint darker brown scales; in some individuals the head can be completely black. Ventrally the base colour is creamish with scattered orange spots. Attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m.

Notes: This species is distributed over a wide area of arid and semi-arid Australia, ranging from western and southern Queensland, western New South Wales, northern South Australia, and central and southern Western Australia. Its principle habitat is open woodland and semi-arid shrubland on red soil plains. An oviparous species, it produces around 20 eggs in a clutch. The main diet comprises lizards but small mammals may also be consumed. This snake is highly venomous and although there is no direct evidence for fatalities resulting from its bite, urgent medical attention should be sought in the event of a bite, because this is most certainly a potentially dangerous species. It is fully protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the SA National Parks and Wildlife Act (1972), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). The conservation status of this species is at present unknown, but it may be considered as potentially vulnerable due to its limited distribution and specialised habitat requirements. Regarded as mostly a very common species over its range, however, some populations may have very restricted distributions.

Longman's Brown Snake ***Pseudonaja carinata* (Longman, 1915)**

Previously known as the 'carinata morph' or the 'banded' form, of *Pseudonaja nuchalis*, recent genetic studies have proven its distinctiveness from that species, necessitating the reinstatement of the original name given it by Heber Longman in 1915 - thus it should now be called *Pseudonaja carinata*.

Diagnosis: This is another medium to large but relatively slender species, Longman's Brown Snake has a small narrow head not distinct from the neck, the eye is large with a round pupil and a pale iris. As presently understood, this species has a distinctive chromosomal morphology (of the $2n=34$ karyotype). Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. The head is brown, the nape is paler with a few darker scales. The base body colour is pale creamish-brown, with the posterior of the body heavily banded with up to 13 (mostly 11) broad blotches or saddle-like bands of black. The dark blotches are slightly narrower than the paler interspaces anteriorly, but are as wide as or slightly wider than the pale areas posteriorly. Within the pale interspaces there are three or four narrow (1 scale wide) dark brown bands. Juveniles have a similar colour and pattern to the adults, with the exception that the head is darker and the broad dark body bands either completely encircle the body or break around the medial area of the ventrals. The WA population appears to have a greater number of bands also, and this may indicate that it is taxonomically distinct in itself. Attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m.

Notes: As presently defined this species is distributed over a wide area of arid and semi-arid Australia, ranging from western and southern Queensland, western New South Wales, possibly north-eastern South Australia, and an apparently isolated population also occurs in the semi-arid south-west Western Australia (although this population has superficial similarities to *Pseudonaja acutirostris*). Its principle habitat is open grassy plains with scattered low shrubs and open woodland. This is an oviparous species, producing up to 20 eggs in a clutch. Feeds mainly on lizards and small mammals.

This snake is highly venomous and although there are no records of fatalities, urgent medical attention should be sought in the event of a bite, because this is most certainly a potentially dangerous species. It is protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the Victorian Wildlife Act (1975)

[but not listed in Schedule 2 of the Victorian Flora and Fauna Guarantee Act (1988)], the SA National Parks and Wildlife Act (1972), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). The conservation status of this species is at present unknown, but it is regarded as mostly a very common species over its range. In some parts of its range however, it may be considered as potentially vulnerable (due to its apparently fragmented distribution and specialised habitat requirements).

Gow's Brown Snake

***Pseudonaja gowi* sp. nov.**

Previously known as the 'southern with black nuchal band morph' of *Pseudonaja nuchalis*, recent genetic studies have proven its specific distinctiveness. As presently understood, this species has a distinctive chromosomal morphology (of the $2n=34$ karyotype). This is another medium to large but relatively slender species with a small narrow head not distinct from the neck, the eye is large with a round pupil and a reddish iris. Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. Very similar in colour and pattern to *Pseudonaja aspidorhyncha*, with its overall plain brown body colour and reduced patterning. However, there is the addition of a broad black band near the neck-nuchal area, and this is occasionally accompanied by a thin secondary collar of black at the anterior end of the neck band. Additionally, there can be a loose vertebral series of black scales on the posterior of the body. The ventral area is creamish with pale orange blotching. Attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m. I hereby choose as Type Locality: Lyndhurst, South Australia and designate the holotype as being the largest specimen of this species from the vicinity of Lyndhurst, SA in the South Australian Museum collection.

Notes: This species has a scattered distribution over arid and semi-arid eastern and central Australia, ranging from south-western Queensland, north-western and far western New South Wales, to eastern and north-eastern South Australia. It is known from a range of temperate semi-arid shrublands. An oviparous species, producing about 20 eggs in a clutch. Feeds mainly on lizards and small mammals. This snake is highly venomous and has almost certainly been the cause of a number of fatalities in the past, so urgent medical attention should be sought in the event of a bite. It is protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the SA National Parks and Wildlife Act (1972) and the Qld Nature Conservation Act (1992). Although regarded as mostly a very common species over its range, some populations may have very restricted distributions. At present, its conservation status is unknown, but this species may be considered as potentially vulnerable in some parts of its range due to its fragmented distribution and specialised habitat requirements. The name 'gowi' honours Australian herpetologist Graeme Francis Gow.

Western Brown Snake ***Pseudonaja imperitor* Wells and Wellington, 1985**

Previously known as the 'Darwin morph' or 'brown with black head' form, of *Pseudonaja nuchalis*, recent genetic studies have proven its distinctiveness from that species, necessitating the re-instatement of the original name given it - thus it should now be called *Pseudonaja imperitor*.

Diagnosis: This is another medium to large but relatively slender snake with a small narrow head not distinct from the neck, the eye is large with a round pupil and a pale iris. As presently understood, this species has a distinctive chromosomal morphology (of the $2n=30$ karyotype). Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. The base colour of the body is uniform tan-brown, and the ventral surface is creamish. The snout is pale creamish-brown, followed by an interocular band of dark brown over the head. The nuchal area can have a few scattered black scales, or occasionally these can form a narrow band over the neck. Juveniles have a similar colour and pattern to the adults, with the exception that the darker interocular area and nape markings are more pronounced. There is quite pronounced seasonal colour change with this form - during the summer the entire body colour becomes much lighter brown, while during the (slightly) cooler winter (or dry season) it changes to a darker brown overall. Attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m. Its distribution is restricted to scattered areas across a wide area of central and northern Australia. *Pseudonaja imperitor* occurs in north-western Queensland, and into much of the Northern Territory, in particular the far north of the NT (including Arnhem Land and some offshore islands) and the far north of Western Australia (in the northern Kimberley region). At present, it is unknown whether this species' distribution is continuous, or merely composed of a few isolated populations. This species occurs in a variety of habitats, but usually open savanna woodland areas with a dense ground cover of grasses are favoured; it has also been found in the vicinity of rock outcroppings. It is an oviparous species, producing less than 20 eggs in a clutch. The main diet is lizards but small mammals may also be consumed. This snake is highly venomous and has caused a number of fatalities, so urgent medical attention should be sought in the event of a bite. Protected under the Territory Parks and Wildlife Conservation Act (1998), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). This is mostly a very common species over its range, however, some populations may have very restricted distributions. As the conservation status of some apparently restricted populations is unknown, they may be considered as potentially vulnerable. The name 'imperitor' means 'commander in chief', and alludes to the defensive behaviour of this species when disturbed.

Kelly's Brown Snake ***Pseudonaja kellyi* Wells and Wellington, 1985**

Previously known as the 'orange with black head morph' or 'hooded' form, of *Pseudonaja nuchalis*, recent genetic studies have proven its distinctiveness from that species, necessitating the re-instatement of the original name given it - thus it should now be called *Pseudonaja kellyi*.

Diagnosis: This is a moderately large but relatively slender species with a small narrow head not distinct from the neck, the eye is large with a round pupil and a pale iris. As presently

understood, this species has a distinctive chromosomal morphology (of the $2n=32$ karyotype). Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. The head and neck region are jet black (or sometimes very dark brown). The base colour of the body is yellowish-orange to orange-brown, with many of the dorsal scales marked with dark brown or black to form a transverse ziz-zag or even a 'herring-bone' pattern over the body. Juveniles have a similar colour and pattern to the adults, with the exception that the snout is paler followed by a darker interocular area and a dark brownish nape. Attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m.

Notes: *Pseudonaja kellyi* is distributed over a wide area of arid and semi-arid Australia, ranging from western and southern Queensland, central-western New South Wales, northern South Australia, the southern half of the Northern Territory, and right across to the far west coast of Western Australia. An apparently isolated population also occurs in the northern Kimberley region as well. It lives in a variety of semi-arid and arid shrubland to open woodland habitats on sand plains and in the vicinity of rocky ranges. This is an oviparous species, and although there are no records of clutch-size its body length indicates that it may producing around 20 eggs in a clutch. The diet comprises lizards and small mammals. This snake is highly venomous and has almost certainly been the cause of a number of fatalities in the past, so urgent medical attention should be sought in the event of a bite. Protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the SA National Parks and Wildlife Act (1972), the Territory Parks and Wildlife Conservation Act (1998), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). The conservation status of *Pseudonaja kellyi* is at present unknown, but some populations may be considered as potentially vulnerable due to its fragmented distribution and specialised habitat requirements. Although regarded as mostly a very common species over its range, some populations appear to have very restricted distributions. The name *kellyi* was bestowed in recognition of the highly defensive behaviour of this snake when confronted - 'kellyi' recalls the Australian folk hero Ned Kelly, who was noted for standing his ground against insurmountable odds.

Mengden's Brown Snake

***Pseudonaja mengdeni* Wells and Wellington, 1985**

Previously known as the 'pale head, grey nape morph' or 'pale-headed' form, of *Pseudonaja nuchalis*, recent genetic studies have proven its distinctiveness from that species, necessitating the re-instatement of the original name given it - thus it should now be called *Pseudonaja mengdeni*.

Diagnosis: This is another medium to large but relatively slender snake with a small narrow head not distinct from the neck, the eye is large with a round pupil and a pale iris. As presently understood, this species has a distinctive chromosomal morphology (of the $2n=34$ karyotype). Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided.

The base body colour can vary from tan-brown through to pale yellow or orange. There is a strong 'herring bone' pattern on the posterior two-thirds of the body caused through the arrangement of darker reddish-brown scales. The head and neck is usually pale brown (snout is paler) with a slightly darker interocular area. The neck is greyish-brown to darker brown, with a dark narrow row of black or very dark brown scales often forming a sharp boundary (sometimes in a 'V' shape) immediately anterior to the nape patch. Juveniles have a similar colour and pattern to the adults, with the exception that the darker interocular area and nape markings are more pronounced. It attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m.

Notes: *Pseudonaja mengdeni* is distributed over a wide area of arid and semi-arid Australia, ranging from central and northern Queensland, north-western New South Wales, most of the Northern Territory, northern and western South Australia, then into adjacent south-eastern Western Australia, and the far west coast of Western Australia from about North-West Cape to Perth. An apparently isolated population also occurs in the vicinity of Broome in north-western WA as well. The principle habitat is open woodland, but it is also known from semi-arid shrubland and grassland on plains country. This is an oviparous species, but nothing is known on its clutch-size, although it could be expected to lay upward of 12 eggs in a clutch. It feeds mainly on lizards and small mammals. This snake is highly venomous and has almost certainly been the cause of a number of fatalities in the past, so urgent medical attention should be sought in the event of a bite. Protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the SA National Parks and Wildlife Act (1972), the Territory Parks and Wildlife Conservation Act (1998), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). Although regarded as mostly a very common species over its range, the conservation status of this species is unknown, and it may be considered as potentially vulnerable in some areas due to its apparently fragmented distribution and specialised habitat requirements. The name 'mengdeni' honours American herpetologist, Gregory Mengden.

Northern Brown Snake ***Pseudonaja nuchalis* Gunther, 1858**

Diagnosis: A large and relatively slender snake, the Northern Brown Snake has a small narrow head not distinct from the neck, large eye size with a round pupil and a pale iris of variable colouration, usually reddish. Based on the original description as well as the available Type Specimen that is consistent with that original description, *Pseudonaja nuchalis* actually represents one of the rarest of the known 'variations' of the complex. In *Pseudonaja nuchalis* sensu stricto the head, neck and throat is jet black. The body has a base colour of yellowish-orange to orange-brown, and there is a series of about 6 broad black bands that do not encircle the body. Each black band has an irregular edge formed by darker brown and black scales, which tends to make the lighter interspaces grade into the darker bands (in other words, not create a clear line of demarcation between the bands). The snout is usually lighter brown, with a slightly darker brown patch on the head; there may be a cluster of brownish scales on the neck also. Ventrally, creamish, with obscure brown edging, becoming darker posteriorly. Some features of this species' scalation are: nasal entire and in contact with preocular, no suboculars, postoculars 2 (occasionally 3), preocular higher than wide and separated from frontal, supralabials 6, infralabials 6, rostral higher than broad, usually extending back onto the top of the snout, temporals 1+2, canthus rostralis very strong, frontal shield longer than wide, and about as wide as a supraocular, body scales smooth in 17 (rarely 19) rows at mid-body, ventrals 180-230, anal divided, and subcaudals 50-70 divided. Attains a maximum size of around 1.5 m in total length, but usually mature specimens are around 1.2 m.

Notes: As presently defined, *Pseudonaja nuchalis* is restricted to the far north of the Northern Territory, centred on Arnhem Land and the adjacent coastal areas. Rather than being the widespread and abundant species it is usually assumed to be, *Pseudonaja nuchalis* as now defined, is probably endangered because of its very restricted known distribution. The principle habitat occupied by this species is open savanna woodland and grassland in association with sandstone outliers of the Arnhem Land escarpment. It is an oviparous species, producing up to 22 eggs in a clutch. The diet is unknown, but it likely feeds mainly on lizards and small mammals. This snake is highly venomous and its bite has likely resulted in several fatalities to date, so urgent medical attention should be sought in the event of a bite. Protected under the Territory Parks and Wildlife Conservation Act (1998). The conservation status of this species is unknown, but this species may be considered as potentially vulnerable or even endangered due to its limited distribution and specialised habitat requirements. The name 'nuchalis' refers to the neck region (nuchal area) and presumably draws attention to the neck colour and pattern.

Placidaserpens gen. nov.

Type Species: *Demansia guttata* Parker, 1926 [New reptiles and a new frog from Queensland. *Annals and Magazine of Natural History*, (9) 17: 665-670]

Diagnosis: As presently defined, a monotypic genus of large snakes in the family Elapidae occurring in Australia and readily identified by the following combination of characters: large robust species with a small head that is barely distinct from the neck; head with moderate canthus rostralis; eyes relatively small with round pupils; iris reddish-yellow, with inner margin thinly bordered with white; nasal and preocular scales in contact; suboculars absent; 1 primary temporal; supralabials 6; infralabials 6; body scales smooth in 19 or 21 rows at mid-body; ventrals 190-220; anal divided; subcaudals 45-70 divided. The maximum size attained is around 1.0 m. in total length, but 0.6 m is an average-sized adult. Populations in the western part of its range (NT) have mainly 19 mid-body rows, while those from the east in central Qld have 21 and these may represent different species. Etymology: The name 'Placidaserpens' means 'peaceful snake', in recognition of the relatively inoffensive nature of some specimens. (but be very careful when handling this species, because they will readily attempt to bite if disturbed). Content: *Placidaserpens guttatus*(Parker, 1926).

Speckled Brown Snake

***Placidaserpens guttatus* (Parker, 1926)**

Diagnosis: A large robust species with a small head that is barely distinct from the neck. The head has a moderate canthus rostralis and the eyes are relatively small with round pupils and a reddish-yellow iris, with a the inner margin thinly bordered with white. The body colour can be almost any shade of brown, but usually tan or orange-brown or yellowish-brown is a common base colouration, with the lower (hidden) edges of each dorsal scale black, and black peppering on the head and neck, with an occasional specimen having a dark brown nuchal blotch. The black-edging of the dorsal scales results in a speckled appearance to the body if the skin is even slightly distended. In another variation, some individuals are strongly marked with up to around 12 black, reddish-brown or dark brown bands or blotches on the body and tail; these markings can vary in width, from being very narrow and barely visible, to being very broad and dominant over the dorsum. Speckling can also occur between the bands as well. Ventrally pale whitish in some, but usually creamish-orange to orange with occasional darker orange spotting or blotching on the ventrals (but this is less distinct posteriorly); the throat and labials are whitish. Juveniles are much lighter than adults and lack any dark colouring to the head and neck. Some significant features of the scalation are: nasal and

preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 19 or 21 rows at mid-body (Populations in the western part of its range (NT) have mainly 19 mid-body rows, while those from the east in central Qld have 21), ventrals 190-220, anal divided, and subcaudals 45-70 divided.

The maximum size attained is around 1.4 m. in total length, although specimens around a metre would be mature; 0.75 m is an average-sized adult.

Notes: Known from a wide area of central and western Queensland and the adjacent north-eastern Northern Territory and north-eastern South Australia. This species inhabits *Astrebla* grasslands with scattered low shrubs on black-soil plains; also found in low rocky hills adjacent to grasslands. Very little has been recorded on the ecology of this abundant species. It is known to be an egg-layer, but the clutch-size is unknown. It feeds mainly on frogs and small lizards and has been known to take small mammals in captivity. This is mainly a diurnal species that shelters in deep earth cracks, particularly in the vicinity of watercourses or ephemeral waterholes. When aroused this nervous snake will raise its head and neck up from the ground, then flatten its neck to form a distinctive cobra-like hood. It is however usually hesitant to bite even if provoked, but extreme care should be shown in any case. This is a highly venomous species and its bite has likely resulted in a number of fatalities, so urgent medical attention should be sought in the event of a bite. Although the conservation status of the species is poorly known, it is protected under the SA National Parks and Wildlife Act (1972), the Territory Parks and Wildlife Conservation Act (1998) and the Qld Nature Conservation Act (1992). It is nevertheless very common within its habitat.

Notopseudonaja gen. nov.

Type Species: *Cacophis modesta* Gunther, 1872 [Seventh account of new species of snakes in the collection of the British Museum. *Annals and Magazine of Natural History*, (4) 9: 13-37].

Diagnosis: As presently defined, a monotypic genus of small snakes in the family Elapidae occurring in Australia, readily identified by the following combination of characters: very small, slender species with a small head that is barely distinct from the neck; weak canthus rostralis when an adult (but not in the juvenile); eyes relatively small with round pupils; iris orange-brown; nasal and preocular scales in contact; suboculars absent; 1 primary temporal; supralabials 6, infralabials 6; body scales smooth in 17 rows at mid-body; ventrals 150-183; anal divided; subcaudals 33-56 divided. Maximum size attained is around 0.6 m. in total length, but 0.45 m is an average-sized adult. Content: *Notopseudonaja modesta* (Gunther, 1872); *Notopseudonaja ramsayi* (Macleay, 1885); *Notopseudonaja sutherlandi* (De Vis, 1884).

Western Ringed Snake

***Notopseudonaja modesta* (Gunther, 1872)**

Diagnosis: A very small, slender species with a small head that is barely distinct from the neck. The head has a weak canthus rostralis when an adult (but not in the juvenile) and the eyes are relatively small with round pupils and an orange-brown iris. The mature body colour may be either light tan, light reddish-brown, yellowish-brown, or greyish (usually) all over, the body mostly being without pattern other than a faint trace of a nuchal band, and a fine speckling of the body caused by some of the dorsals being marked with paler bases and slightly darker centres. Juveniles and immatures however, are brightly coloured (usually reddish or orange-brown) with about 4 to 12 bold narrow black transverse bands. The top of the head has a black patch which extends down the side of the head to include the eyes and part of the supralabials, and there is another black patch on the nape. Ventrally, usually

creamish or white with occasional specimens flecked with orange on the ventrals. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 17 rows at mid-body, ventrals 150-183, anal divided, and subcaudals 33-56 divided. The maximum size attained is around 0.6 m. in total length, but 0.45 m is an average-sized adult.

Notes: Found over a large part of arid and semi-arid north-western and central Western Australia and adjacent north-western Northern Territory. Occurs in a wide range of arid and semi-arid habitats from gibber deserts, sand deserts, tropical savanna, open woodland, shrubland, sandplains, and rocky ranges. Usually found in association with sandy soils in densely vegetated shrublands and open woodlands with *Triodia* grass ground cover. This is an oviparous species, producing up to 11 eggs in a clutch. Feeds mainly on lizards. Although this is mainly a diurnal species it may also be encountered during warm evenings. They may be found in the burrows of other animals or beneath or associated with low vegetation, such as *Triodia* tussocks and deep litter under shrubs. They can also be found active at night during warm weather. When aroused, it will raise its head from the ground and place the neck in an 'S' shape, while hissing loudly. Its small size has led to the belief that this is virtually an innocuous species. Although this is reportedly only a moderately venomous species and although its bite has not resulted in any fatalities to date, I believe that urgent medical attention should be sought in the event of a bite, particularly from a large specimen. It would be sensible to exert caution when handling this snake until the nature of its venom is better known. Although usually inoffensive, even rather placid despite being moderately disturbed, they will try to bite if given the opportunity. Protected under the Territory Parks and Wildlife Conservation Act (1998) and the WA Wildlife Conservation Act 1950 (as amended). Regarded as common. The name 'modesta' means 'modest', and presumably refers to the plain colouration and patterning of the adult of this species.

Southern Ringed Snake ***Notopseudonaja ramsayi* (Macleay, 1885)**

Diagnosis: A very small, slender species with a small head that is barely distinct from the neck. The head has a weak canthus rostralis when an adult (but not in the juvenile) and the eyes are relatively small with round pupils and an orange-brown iris. The mature body colour may be either light tan, light reddish-brown, yellowish-brown, or greyish (usually) all over, the body mostly being without pattern other than a faint trace of a nuchal band, and a fine speckling of the body caused by some of the dorsals being marked with paler bases and slightly darker centres. Juveniles and immatures however, are brightly coloured (usually reddish or orange-brown) with about 4 to 12 bold narrow black transverse bands. The top of the head has a black patch which extends down the side of the head to include the eyes and part of the supralabials, and there is another black patch on the nape. Ventrally, usually creamish or white with occasional specimens flecked with orange on the ventrals. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 17 rows at mid-body, ventrals 150-183, anal divided, and subcaudals 33-56 divided. The maximum size attained is only around 0.5 m. in total length, but 0.3 m is an average-sized adult.

Notes: As herein defined it is found over much of arid and semi-arid eastern and southern Australia, ranging from far western New South Wales, south-western Queensland, through most of central and northern South Australia, most of the southern part of the Northern Territory and into south-eastern Western Australia. Occurs across a wide range of arid and semi-arid habitats from gibber deserts, sand deserts, sub-tropical woodland, temperate semi-arid open woodland, shrubland, sandplains, and rocky ranges. Usually found in association with sandy soils in densely vegetated shrublands and open woodlands with *Triodia* or

Plectrachne grass ground cover. Oviparous, producing about 6 eggs in a clutch. Feeds mainly on lizards. This is mainly a diurnal species that seeks shelter in lizard burrows beneath or associated with low vegetation. They can also be found active at night during warm weather. When aroused, it will raise its head from the ground and place the neck in an 'S' shape, while hissing loudly. As in the case of *N. modesta* caution should be exerted in the case of any bite from this species. Its small size has led to the belief that this is virtually an innocuous species, and its overall inoffensive behaviour makes one think that it is virtually harmless. However, lizards are quickly subdued with its venom and I think that it would be wise to seek urgent medical attention in the event of a bite - particularly from a large specimen. It would be sensible to exert caution when handling this snake until the nature of its venom is better known. Protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the SA National Parks and Wildlife Act (1972), the Territory Parks and Wildlife Conservation Act (1998), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). Regarded as common. The name 'ramsayi' honours 19th century Australian herpetologist Edward Pierson Ramsay.

Northern Ringed Snake ***Notopseudonaja sutherlandi* (De Vis, 1884)**

Diagnosis: This is another very small, slender species with a small head that is barely distinct from the neck. It has been in the past regarded as a synonym of *Pseudonaja nuchalis* by Cogger et al (1983) but this is now known to be incorrect (see Mengden, 1985 who showed that it was actually part of the *modesta* group). The head has a weak canthus rostralis when an adult (but not in the juvenile) and the eyes are relatively small with round pupils and an orange-brown iris. The mature body colour may be either light tan, light reddish-brown, yellowish-brown, or greyish (usually) all over, the body mostly being without pattern other than a faint trace of a nuchal band, and a fine speckling of the body caused by some of the dorsals being marked with paler bases and slightly darker centres. Juveniles and immatures however, are brightly coloured (usually reddish or orange-brown) with about 4 to 12 bold narrow black transverse bands. The top of the head has a black patch which extends down the side of the head to include the eyes and part of the supralabials, and there is another black patch on the nape. Ventrally, usually creamish or white with occasional specimens flecked with orange on the ventrals. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 17 rows at mid-body, ventrals 150-183, anal divided, and subcaudals 33-56 divided. The maximum size attained is only around 0.45 m.

Notes: Found over a wide area of semi-arid tropical northern Australia, ranging from eastern central Northern Territory and most of northern Queensland. It occurs across a wide range of semi-arid habitats from tropical savanna, open woodland, and shrubland. Usually found in association with sandy soils in densely vegetated shrublands and open woodlands. This species is oviparous in its reproductive habits, and feeds only on lizards. This is mainly a diurnal species that seeks shelter in earth cracks, or beneath or associated with low vegetation and ground litter. They can also be found active at night during warm weather. When aroused, it will raise its head from the ground and place the neck in an 'S' shape, while hissing loudly. Its small size has led to the belief that this is virtually an innocuous species, and its overall inoffensive behaviour makes one think that it is virtually harmless. However, I think that it would be wise to seek urgent medical attention in the event of a bite - particularly from a large specimen. It would be sensible to exert caution when handling this snake until the nature of its venom is better known. Although usually inoffensive, even rather placid despite being

moderately disturbed, they will try to bite if given the opportunity. Protected under the Territory Parks and Wildlife Conservation Act (1998) and the Qld Nature Conservation Act (1992). Regarded as common.

Dugitophis gen. nov.

Type Species: *Pseudonaja affinis* Gunther, 1872 [Seventh account of new species of snakes in the collection of the British Museum. *Annals and Magazine of Natural History*, (4) 9: 13-37].

Diagnosis: A genus of medium to large elongate snakes of the family Elapidae occurring in south-western Australia, and readily identified by the following combination of characters: medium-sized to large and slender species with a small head that is not distinct from the neck; head with strong canthus rostralis; eyes relatively small with round pupils and a blackish iris, with a thin yellow or orange ring around the pupil; nasal and preocular scales in contact; suboculars absent; 1 primary temporal; supralabials 6; infralabials 6; body scales smooth in 19 (rarely 17 or 21) rows at mid-body; ventrals 190-230; anal divided; and, subcaudals 50-70 divided. Content: *Dugitophis affinis affinis* (Gunther, 1872); *Dugitophis affinis exilis* (Storr, 1989); *Dugitophis affinis tanneri* (Worrell, 1961).

Dugite

***Dugitophis affinis affinis* (Gunther, 1872)**

Diagnosis: This is a medium-sized and slender species with a small head that is not distinct from the neck. The head has a strong canthus rostralis and the eyes are relatively small with round pupils and a blackish iris, with a thin yellow or orange ring around the pupil. The body colour can be any shade of brown, but usually dark olive or greyish-brown is a common base colouration. Some of the dorsal scales are coloured black, and these black scales can be sparsely scattered over the dorsum, or clustered together in patches without any particular pattern being created. The blotched form (known as the "Kabada") can be so heavily marked that it is almost black with only scattered sections of a lighter tan-yellow base colour evident, while in others the dark blotching is more evenly distributed over the body. Ventrally, usually whitish with incomplete edging on the margins of the ventral scales. Juveniles usually have a black patch on the top of the head and another on the neck, with a greenish-brown to light yellowish body colour, and a faint darker herring-bone pattern overlying the base colour, and a white belly with scattered brownish-red spotting. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 19 (rarely 17 or 21) rows at mid-body, ventrals 190-230, anal divided, and subcaudals 50-70 divided. The maximum size attained is around 2.0 m in total length, but 1.2 m is an average-sized adult.

Notes: Known only from the south-western and southern coastal parts of Western Australia, and adjacent south-west of South Australia. This species occurs in a wide variety of vegetation communities, ranging from wet sclerophyll forest, rocky areas, coastal dune shrub-heath, semi-arid woodland, usually on sandy soils. It is known to be an egg-laying species, producing up to 30 in a clutch (sometimes two clutches are laid in a season) between November and February. Eggs may hatch between February and May, and juveniles can be around 200 mm at hatching. Feeds mainly on lizards and small mammals, but will also eat other snakes on occasions. This is a diurnal and terrestrial species that shelters in holes in the ground or under logs or rocks. When aroused, it often raises its head from the ground and places the neck in an 'S' shape, while hissing loudly. Sometimes this species is quite inoffensive and can be rather placid even when moderately disturbed. However, on other

occasions it will not hesitate to bite if given the opportunity, so extreme care should be shown. This is a highly venomous species and its bite has resulted in a number of fatalities, so urgent medical attention should be sought in the event of a bite. Protected under the SA National Parks and Wildlife Act (1972) and the WA Wildlife Conservation Act 1950 (as amended). Common wherever it occurs. The name 'affinis' means 'related to', and was bestowed because of its alleged similarity in appearance with *Pseudonaja nuchalis*.

Rottnest Island Dugite
***Dugitophis affinis exilis* (Storr, 1989)**

Diagnosis: This is a smaller and more slender form than the nominate subspecies. It has a small head that is not distinct from the neck, with a medium canthus rostralis. It has small eyes with round pupils and a blackish iris, with a thin yellow or orange ring around the pupil. The colour of the dorsal and ventral areas is black. Juveniles from Rottnest Island are very similar to the juveniles of the nominate form. They usually have a black patch on the top of the head and another on the neck, with a light brownish to yellowish body colour, and a faint darker herring-bone pattern overlying the base colour, and a white belly with scattered brownish-red spotting. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 19 (rarely 17 or 21) rows at mid-body, ventrals 190-230, anal divided, and subcaudals 50-70 divided. The maximum size attained is around 1.0 m. in total length, but 0.6 m is an average-sized adult.

Notes: Restricted to Rottnest Island in south-western Western Australia, where it occupies semi-arid temperate shrublands and heath communities on sandy soils. Its reproductive biology is poorly known, although it is egg-laying, producing up to 20 in a clutch. The diet comprises mainly small lizards but small mammals may also be taken. This is a diurnal and terrestrial species that shelters in holes in the ground or under rocks. When aroused, it will raise its head from the ground and place the neck in an 'S' shape, while hissing loudly. This snake is usually inoffensive, but they will not hesitate to bite if given the opportunity, so extreme care should be shown. It is a highly venomous snake and although there are no records of fatalities, urgent medical attention should be sought in the event of a bite, because this is most certainly a potentially dangerous subspecies. The nominate subspecies has caused a number of fatalities. Protected under the WA Wildlife Conservation Act 1950 (as amended). Status unknown, but this subspecies may be considered as potentially vulnerable due to its limited distribution and specialised habitat requirements.

Tanner's Brown Snake
***Dugitophis affinis tanneri* (Worrell, 1961)**

Diagnosis: This race is also smaller and more slender than the nominate subspecies. It has a small head that is not distinct from the neck, with a medium canthus rostralis. It has small eyes with round pupils and a blackish iris, with a thin yellow or orange ring around the pupil. The body colour is usually dark chestnut-brown without spotting. Ventrally, usually whitish with incomplete edging on the margins of the ventral scales. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 19 (rarely 17 or 21) rows at mid-body, ventrals 190-230, anal divided, and subcaudals 50-70 divided. The maximum size attained is around 2.0 m. in total length, but 1.0 m is an average-sized specimen.

Notes: Restricted to two of the islands of the Recherche Archipelago in south-western Western Australia (Boxer Island and Figure-of-Eight Island), where it occupies semi-arid temperate low woodland and shrub-heath communities with limestone on sandy soils. It is

believed to produce up to 20 eggs in a clutch, and feed mainly on small lizards. This is a diurnal and terrestrial subspecies that shelters in holes in the ground or under rocks. When aroused, it will raise its head from the ground and place the neck in an 'S' shape, while hissing loudly. As the nominate subspecies has caused a number of fatalities, and the species overall is highly venomous, urgent medical attention should be sought in the event of a bite from this particular subspecies. Protected under the WA Wildlife Conservation Act 1950 (as amended). Status unknown, but this subspecies may be considered as potentially vulnerable due to its limited distribution and specialised habitat requirements. The name 'tanneri' honours Australian herpetologist Charles Tanner (1911-1996)

Genus Euprepiosoma Fitzinger, 1860

Peninsula Brown Snake

Euprepiosoma infracula (Waite, 1925)

Diagnosis: This is a large but slender species with a small head that is not distinct from the neck. The head has a strong canthus rostralis and the eyes are relatively small with round pupils and a pale brown iris. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal, supralabials 6, infralabials 6, body scales smooth in 17 rows at mid-body, ventrals 185-235, anal divided, and subcaudals 45-75 divided. The body colour can be any shade of pale or rich brown, even black, but usually dark brown or greyish-brown are common base colourations. Sometimes pale yellowish specimens may be found also. Some of the dorsal scales are coloured black, and these black scales may be densely scattered over the dorsum with the spotting being often concentrated along the vertebral line. Occasionally distinctly banded specimens occur in some areas. The head can be flecked or spotted with black, and usually has a shiny or glossy appearance. Ventrally, pale brown with the centre of the belly more towards dark grey with black spots, or in some specimens a greyish mottling pattern may dominate. The maximum size attained is around 1.5 m. in total length, but an average adult is usually about 1.2m.

Notes: Restricted to a small area of coastal southern South Australia, principally Eyre Peninsula, York Peninsula and a few nearby islands, where it inhabits a range of quite different habitats, ranging from areas of temperate semi-arid heath and shrubland on sandy soil or dunes, to dry sclerophyll woodland. Known to also inhabit limestone outcroppings in some areas also. Surprisingly little biological data is available on this very common species. It is known to be an egg-laying species, producing up to 20 in a clutch. Peninsula Brown Snakes feed mainly on small lizards, ground-nesting birds, frogs and small mammals. It is diurnal and very fast-moving, and is often observed basking on roads - consequently, many are killed by traffic. Although a rather inoffensive and shy species, it is still highly venomous. There are few records of bites, and symptoms may be severe, so urgent medical attention should always be sought in the event of a bite, because this is most certainly a potentially dangerous species. Protected under the SA National Parks and Wildlife Act (1972). Very common within its habitat, despite its restricted distribution.

Ingram's Brown Snake

Euprepiosoma ingrami (Boulenger, 1908)

Diagnosis: This is a large and robust species with a small head that is not distinct from the neck. The head has a strong canthus rostralis and the eyes are relatively small with round pupils and a dull orange-brown (almost blackish) iris. Some significant features of the scalation are: nasal and preocular scales in contact, suboculars absent, 1 primary temporal,

supralabials 6, infralabials 7, body scales smooth in 17 rows at mid-body, ventrals 190-230, anal divided, and subcaudals 50-70 divided. The body colour can be a variety of browns, ranging from very dark glossy brown (almost blackish-brown), to a form that is a light olive-brown, or even a golden brown overall. Another variation can be dark brown anteriorly and golden brown posteriorly, while another can be a pale to rich yellowish-brown, with the head and nape being dark brown to black. The dorsal scales are always tipped with dark brown. Ventrally, this species may be either pale yellow, bright yellow or even orange, with bright orange spots near the edges of the ventral scales; the chin and labials are creamish. The buccal cavity colour is blackish. The maximum size attained is around 2.0 m. in total length, but 1.7 m is a large adult.

Notes: Restricted to a small area of north-western Queensland and the adjacent north-eastern Northern Territory, and an apparently isolated population in the Kimberley region of far north-east Western Australia (in the vicinity of Kununurra). Generally, it inhabits semi-arid black-soil plains with *Astrebla* grass cover and scattered low shrubs. Its favoured habitat is usually subject to seasonal inundation by late summer rains. Oviparous, producing up to 12 eggs in a clutch. Juveniles feed mainly on lizards, and adults on small mammals such as the Long-haired Rat (*Rattus villosissimus*). A secretive, mainly diurnal snake that may be encountered basking or active in the early morning. As the temperature increases, it will seek shelter in deep earth cracks until cooler afternoon conditions when it may emerge again. Although this species has been seldom observed in the field, it is more often found following heavy rain when the ground is saturated and they are forced to the surface. It may also be occasionally found active at night during hot weather. When aroused this snake will raise its head and neck up from the ground, then flatten its neck to form a distinctive cobra-like hood in a similar manner to a Speckled Brown Snake. It is usually hesitant to bite even if provoked, but extreme care should be shown in any case. This snake is highly venomous and although its bite has not resulted in any fatalities to date, urgent medical attention should be sought in the event of a bite, because this is most certainly a potentially dangerous species. Protected under the Territory Parks and Wildlife Conservation Act (1998), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). Seldom found, but known to be common within its habitat. Named for the English ornithologist Collingwood Ingram.

Eastern Brown Snake

***Euprepisoma textilis* (Dumeril, Bibron and Dumeril, 1854)**

Diagnosis: Variation in morphology suggests that this species may be composite. This is a very large and robust species with a small head that is not distinct from the neck. The head has a strong canthus rostralis and the eyes are relatively small with round pupils and a dull pale brown (to almost blackish) iris with a broken yellow or orange ring around the pupil. Some significant features of the scalation are: frontal straight-sided and longer than wide (and about as wide as a supraocular), nasal and preocular scales in contact, preocular higher than wide, postoculars 2, suboculars absent, temporals 1+2 (ie only 1 primary temporal), supralabials 6, infralabials 6, body scales smooth in 17 rows at mid-body, ventrals 185-235, anal divided, and subcaudals 45-76 divided. As presently defined, this is a highly variable species with a confusing array of colour and pattern combinations. The base body colour can be any of variety of browns or paler colours, ranging from greyish-brown, olive-brown, reddish-brown, orange or golden brown and very dark (almost blackish-) brown, even black. In very dark specimens the head and nape may be much paler. In some specimens, the dorsal scales are faintly edged with dark grey, but usually the (adult) body is unpatterned. In some populations there can be seasonal variation in tonal colouration, with specimens tending to be lighter during the Summer months and greyish-black during the late Autumn to early Spring

period. Although the body is usually unpatterned in mature specimens, in juveniles and even immatures up to around 1 m. in length, can have a series of narrow transverse darker bands, that are usually most distinct in the juveniles but barely noticeable by the time the snake reaches maturity. In some populations the juveniles lack this narrow banding, and just have a black patch on the top of the head (but the snout is pale) and another black patch on the nape - with the body usually a uniform light olive-brown. Ventrally, this species may be either pale yellow or creamish, with bright orange spots or greyish blotches on the ventral scales. The buccal cavity colour is pinkish. The maximum size attained is around 2.3 m. in total length, but 1.8 m is a large adult.

Notes: As its common name suggests, this species occurs mainly in eastern Australia, ranging from north Queensland, south to most of eastern, central and south western New South Wales, and the Australian Capital Territory, throughout Victoria and into the south-east of South Australia. However, a number of outlying populations are also known from central Australia (southern and central parts of the Northern Territory) and from the southeast Kimberley region of northern Western Australia. It is also known from New Guinea. More often found in woodland habitats on sandy or loamy soils with a grassy and shrub understorey, and often in or adjacent to rocky ranges. Known from a wide range of other habitats, including wet and dry sclerophyll forest, coastal dune heath and shrublands, open grassy floodplains, and agricultural habitats. This is an oviparous species, producing up to 35 eggs in a clutch, which take about 80 days to hatch. Feeds mainly on lizards, ground-nesting birds, and small mammals, but will also eat other snakes. This is mainly a diurnal, highly alert and swift-moving snake of drier habitats. In warmer weather they can occasionally be found active on the ground at night. They shelter in earth cracks, mammal burrows, beneath logs or stumps, inside hollow logs and beneath rocks, indeed almost any object laying flush with the ground. In agricultural areas they readily occupy hay stacks and other farm produce and equipment, and this could partly explain the distinct distribution pattern of the species. Over the years it is possible that this species has been accidentally transported all over Australia (and into Papua-New Guinea?) in shipments of produce, building materials and even in farm machinery. However some regard the New Guinea population as a distinct species in its own right. This species can be highly defensive in its behaviour if disturbed, rapidly striking and delivering several bites before attempting to escape. When aroused, it often raises its head from the ground, opens its mouth in a chewing action and places the neck in an 'S' shape, prior to delivering a strike. Occasionally, specimens may raise the head and fore-body from the ground and just flatten the neck cobra-like, but this is a much less dangerous posture than the 'S'-shape. People have been often bitten by this snake while trying to kill it, as most underestimate how quick this species can move if the need arises. This snake is highly venomous and its bite has resulted in many fatalities to date, so urgent medical attention should be sought in the event of a bite. Protected under the New South Wales National Parks and Wildlife Act (1974) but not listed in that State as a Threatened Species in any of the Schedules of the NSW Threatened Species Conservation Act (1995). Also protected under the ACT Nature Conservation Act (1980), the Victorian Wildlife Act (1975) [but not listed in Schedule 2 of the Victorian Flora and Fauna Guarantee Act (1988)], the SA National Parks and Wildlife Act (1972), the Territory Parks and Wildlife Conservation Act (1998), the WA Wildlife Conservation Act 1950 (as amended) and the Qld Nature Conservation Act (1992). Regarded as common throughout most of its distribution. The name 'textilis' means 'woven' as in textiles, and probably refers to the regular body scalation of the species.

References

- Agar, N.S., Board, P.G., Gruca, M.A. and Shine, R. 1977 Studies on the blood of Australian Elapid snakes. 2. Red cell enzymes and glycolytic intermediates. *Comparative Biochemistry and Physiology*, (B) 56: 357-360
- Annable, T. 1986 Subcaudal scalation analysis of *Pseudonaja textilis* (Dumeril and Bibron) in the eastern Riverina region. *Herpetofauna*, 16 (2): 40-42 [not 1985 as printed on cover]
- Anonymous 1874 Snake poisoning by the Indian and Australian reptiles, and the ammonia treatment. *Lancet*, (1874) 2: 418
- Anonymous 1920 Tom Morrisey, Master of Snakes. A remarkable Australian character who knows neither pain nor fear. *Capricornian*, 24 July
- Anonymous 1929 Australian snakes. *Sydney Morning Herald*, 2 November: 12
- Anonymous 1932 Baby bitten by Brown Snake. *Sydney Morning Herald*, 2 November: 18
- Anonymous 1952 Australian snakes. Bites and treatment (First Aid) 1952. Australian Museum, School Leaflet No 3: 4 pages
- Anonymous 1953 Experts differ on Brown Snakes, Taipans. *Courier Mail* [Brisbane], January
- Anonymous 1954 Know Australia. Snakes. *A.M. Weekly* (Sydney), 19 October
- Anonymous 1963 Australian snakes. Australian Museum, Sydney [Leaflet No 59: pp. 1-10]
- Anonymous 1967 'Stick' was a Brown Snake. *Daily Telegraph* [Sydney], 4 December
- Anonymous 1968 Australia's venomous snakes, the distribution and identification of the more important deadly snakes and treatment of envenomation. [Pp. 39-82]. In: Garnet, J.R. (Editor): *Venomous Australian Animals Dangerous to Man*. Commonwealth Serum Laboratories, Parkville [Pp. 1-86 + insert]
- Anonymous 1968 [Photograph of Australian Brown Snake.] *Geographical Magazine*, 6: 1243
- Anonymous 1993 Serendipity strikes again, or Herpetologists rediscover the lost Pigmy Blue-tongue inside a dead Brown Snake. *On the Brink ! Newsletter of the Endangered Species Program*, No 3: 1
- Banks, C.B. 1981 Notes on seasonal colour changes in a Western Brown Snake. *Herpetofauna*, 13: 29-30
- Banks, C.B. 1983 Reproduction in two species of captive Brown Snakes, genus *Pseudonaja*. *Herpetological Review*, 14 (3): 77-79
- Barnett, D., Howden, M.E.H. and Spence, I. 1980 A neurotoxin of novel structural type from the venom of the Australian Common Brown Snake. *Naturwissenschaften*, 67 (8): 405-406

- Barrett, C.L. 1950 Reptiles of Australia: Crocodiles, Snakes and Lizards. Cassell, Sydney [Pp. i-xiii, 1-168]
- Baverstock, P.R. and Schwaner, T.D. 1985 Phylogeny of Australian Elapid snakes: The genetic approach. [Pp. 159-164]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): Biology of Australasian Frogs and Reptiles. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Berncastle, J. 1870 On the distinction between the harmless and venomous snakes of Australia. Australian Medical Gazette, 2: 21-22
- Beste, H. 1970 Reptiles of the Hattah district, Victoria. Victorian Naturalist, 87: 262-265
- Bland, W. 1861 Bites of the venomous snakes of Australia. Australian Medical Journal, 6: 1-6
- Board, P.G., Roberts, J. and Shine, R. 1977 Studies on the blood of Australian Elapid snakes. I. Morphology and composition. Comparative Biochemistry and Physiology, (B) 56: 353-356
- Boulenger, G.A. 1896 Catalogue of the Snakes in the British Museum. Volume 3. British Museum (Natural History), London
- Boulenger, G.A. 1908 Description of a new Elapine snake from Australia. Annals and Magazine of Natural History, (8) 1: 333-334
- Bradshaw, S.D. 1977 Reptiles and their adaptation to arid environments. [Pp. 145-160]. In: Messel, H. and Butler, S.T. (Editors): Australian Animals and their Environment. Shakespeare Head Press, Sydney
- Breeden, S. and Breeden, K. 1967 Animals of Eastern Australia. A Photographic Account of the Mammals, Reptiles and Amphibia. Sydney [Pp. 1-128]
- Broad, A.J., Sutherland, S.K. and Coulter, A.R. 1979 The lethality in mice of dangerous Australian and other snake venoms. Toxicon, 17: 664-667
- Brooker, M. and Wombey, J. 1986 Some observations on the herpetofauna of the Macquarie Marshes region, NSW, with special reference to Chelidae. Australian Zoologist, 23 (1): 1-4
- Burbidge, A.A. and Wyre, G. 1993 Conservation of reptiles and frogs in Western Australia. [Pp. 43-47]. In: Lunney, D. and Ayers, D. (Editors): Herpetology in Australia: A diverse discipline. Royal Zoological Society of New South Wales, Sydney [This issue represented an unnumbered "Transactions of the Royal Zoological Society of New South Wales"]
- Bush, B. 1981 Reptiles of the Kalgoorlie-Esperance Region. Vanguard Press, Perth
- Bush, B. 1989 Ontogenetic colour change in the Gwarder, *Pseudonaja nuchalis*. Western Australian Naturalist, 18 (2): 25
- Bush, B. 1990 Polymorphism in captive bred siblings of the snake, *Pseudonaja nuchalis*. Herpetofauna, 19 (2): 28-32 [what!?!]

- Bush, B., Maryan, B., Browne-Cooper, R. and Robinson, D. 1995 Reptiles and Frogs of the Perth Region. University of Western Australia Press, Perth [Pp. i-xiv. 1-226]
- Bustard, H.R. 1968 The reptiles of Merriwindi State Forest, Pilliga West, northern New South Wales, Australia. *Herpetologica*, 24 (2): 131-140
- Bustard, H.R. 1968 *Pygopus nigriceps* (Fischer): A lizard mimicking a venomous snake. *British Journal of Herpetology*, 4 (2): 22-24
- Calmette, A. 1898 On the curative value of the 'Antivenomous Serum' in relation to the poisons of Australian and Indian snakes. *Intercolonial Medical Journal of Australasia*, 3: 192-197
- Campbell, C.H. 1976 Snake Bite, Snake Venoms and Venomous Snakes of Australia and New Guinea. An annotated bibliography. School of Public Health and Tropical Medicine, University of Sydney, Service Publication No 13: i-ix, 1-222
- Cann, J. 1986 *Snakes Alive*. Kangaroo Press, Sydney
- Challen, R.G. 1975 Venomous Australian snakes and antivenenes. *Australian Journal of Hospital Pharmacy*, 5 (1): 4-16
- Chapman, A. and Dell, J. 1985 Biology and zoogeography of the amphibians and reptiles of the Western Australian wheatbelt. *Records of the Western Australian Museum*, 12 (1): 1-46
- Chester, A. and Crawford, G.P.M. 1980 Envenomation by a juvenile Gwardar. *Medical Journal of Australia*, 1980 (2): 462
- Cogger, H.G. 1965 Reptiles and frogs of Australia's arid regions. *Australian Natural History*, 15 (4): 128-131
- Cogger, H.G. 1965 Reptiles and Amphibians. [Pp. 185-187]. In: *Australian Encyclopaedia Yearbook*. Grolier Society, Sydney
- Cogger, H.G. 1967 *Australian Reptiles in Colour*. Reed, Sydney [Pp. 1-112]
- Cogger, H.G. 1967 The snakes of Australia. [Pp. 152-158]. In: McMichael, D.F. (Editor): *Treasury of Australian Wildlife*. Ure Smith, Sydney
- Cogger, H.G. 1971 The venomous snakes of Australia and Melanesia. [Pp. 35-77]. In: Bucherl, W. and Buckley, E. (Editors): *Venomous Animals and Their Venoms*. Volume 2. *Venomous Vertebrates*. Academic Press, New York
- Cogger, H.G. 1972 *Australian Reptiles in Colour*. Reed, Sydney ['Revised' Edition; Pp. 1-112]
- Cogger, H.G. 1975 *Reptiles and Amphibians of Australia*. Reed, Sydney [1st Edition]
- Cogger, H.G. 1979 Type specimens of reptiles and amphibians in the Australian Museum.. *Records of the Australian Museum*, 32: 164-210

- Cogger, H.G. 1979 Reptiles and Amphibians of Australia. Reed, Sydney [2nd Edition]
- Cogger, H.G. 1983 Reptiles and Amphibians of Australia. Reed, Sydney [3rd Edition]
- Cogger, H.G. 1984 Reptiles in the Australian arid zone. [Pp. 235-252]. In: Cogger, H.G. and Cameron, E.E. (Editors): Arid Australia. Australian Museum, Sydney
- Cogger, H.G. 1985 Australian Proteroglyphous snakes - An historical overview. [Pp. 143-154]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): Biology of Australasian Frogs and Reptiles. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Cogger, H.G. 1986 Reptiles and Amphibians of Australia. Reed, Sydney [4th Edition; Pp. 1-688]
- Cogger, H.G. 1986 Comments on the biogeography of the Elapid snakes of southeastern Australia. [Pp. 16-19]. In: Longmore, R. (Editor) 1986): Atlas of Elapid snakes of Australia. Bureau of Flora and Fauna, Canberra [Australian Flora and Fauna Series, Number 7]
- Cogger, H.G. 1986 Reptiles and Amphibians of Australia. Reed, Sydney [4th Edition; Pp. 1-688]
- Cogger, H.G. 1987 The venomous land snakes. [Pp. 340-355]. In: Covacevich, J., Davie, P. and Pearn, J. (Editors): Toxic Plants & Animals. A Guide for Australia. Queensland Museum, Brisbane
- Cogger, H.G. 1988 Reptiles and Amphibians of Australia. Reed, Sydney [A reprinting, but due to changes, should be regarded as a 5th Edition]
- Cogger, H.G. 1992 Reptiles and Amphibians of Australia. Reed, Sydney [6th Edition]
- Cogger, H.G. 1993 General description and definition of the Order Squamata. [Pp. 157-171]. In: Glasby, C.J., Ross, G.J.B. and Beesley, P.L. (Editors): Fauna of Australia. Volume 2A. Amphibia and Reptilia. Australian Government Publishing Service, Canberra
- Cogger, H.G. 1994 Reptiles and Amphibians of Australia. Reed, Sydney [7th Edition - another substantially altered edition that included an updated Appendix of recently described taxa]
- Cogger, H.G. 1996 Reptiles and Amphibians of Australia. Reed, Sydney [8th Edition]
- Cogger, H.G. 2000 Reptiles and Amphibians of Australia. New Holland, Sydney [9th Edition]
- Cogger, H.G. 2001 State of biodiversity. Conserving Australia's reptiles. Are we serious? Life Line, Bulletin of the Community Biodiversity Network [Sydney], 7 (2): 23-26
- Cogger, H.G. and Heatwole, H.F. 1981 The Australian reptiles: Origins, biogeography, distribution patterns and island evolution. [Pp. 1331-1373] In: Keast, A. (Editor): Ecological Biogeography of Australia. Junk, The Hague [Monographiae Biologicae, Volume 41]

Cogger, H.G. and Heatwole, H.F. 1984 The Australian reptiles: Origins, biogeography, distribution patterns and island evolution. [Pp. 343-370]. In: Archer, M.A. and Clayton, G. (Editors): Vertebrate Zoogeography and Evolution in Australia. (Animals in Space Time). Hesperian Press, Carlisle [Republication of Cogger, H.G. and Heatwole, H. (1981)]

Cogger, H.G. and Lindner, D.A. 1974 Frogs and reptiles. [Pp. 63-107] In: Frith, H.J. and Calaby, J.H. (Editors): Fauna Survey of the Port Essington District, Coburg Peninsula, Northern Territory of Australia. CSIRO Division of Wildlife Research, Technical Paper No 28

Cogger, H.G. and Webber, P. 1976 Reptiles and Amphibians. [Pp. 67-84]. In: Broadbent, J. and Clark, S. (Editors): A Faunal Survey of East Australian Rainforests, Interim Report. Australian Museum, Sydney

Cogger, H.G., Cameron, E.E. and Cogger, H.M. 1983 Zoological Catalogue of Australia. Volume 1. Amphibia and Reptilia. Australian Government Publishing Service, Canberra [Pp. i-vi, 1-313]

Cogger, H.G., Cameron, E.E., Sadler, R.A. and Egger, P. 1993 The Action Plan for Australian Reptiles. Australian Nature Conservation Agency [Canberra], Endangered Species Program, Project No 124 [Pp. 1-254]

Coulter, A.R., Broad, A.J. and Sutherland, S.K. 1979 Isolation and properties of a high molecular weight neurotoxin from the Eastern Brown Snake (*Pseudonaja textilis*). In: Chubb, I.W. and Greffen, L.B. (Editors): Neurotoxins: Fundamental and Clinical Advances. Adelaide University Union Press, Adelaide

Covacevich, J.A. 1970 The Snakes of Brisbane. Queensland Museum, Brisbane

Covacevich, J.A. 1976 Amphibians and Reptiles. [Pp. 16-18]. In: Covacevich, J. (Editor): Fauna of Eastern Australian Rainforests - Preliminary report on sites surveyed by the Queensland Museum in mid-eastern and north-eastern Queensland. Queensland Museum, Brisbane [March 1976]

Covacevich, J.A. 1980 Medically significant terrestrial snakes of northern Australia. In: Blackall, M.I. and McDonald, H. (Editors): Child Health, Past, Present and Future. Proceedings of the Royal Children's Hospital Academic Week, Brisbane. Royal Children's Hospital, Brisbane

Covacevich, J.A. 1981 On the resemblance between the Western Brown Snake and the Small-scaled Snake. *Herpetofauna*, 12 (2): 30

Covacevich, J.A. 1986 Aspects of the biogeography of the Elapid snakes of northeastern Australia. [Pp. 20-24]. In: Longmore, R. (Editor) (1986): Atlas of Elapid Snakes of Australia. Bureau of Flora and Fauna, Canberra [Australian Flora and Fauna Series Number 7]

Covacevich, J.A. 1987 Reptiles of far northern Cape York Peninsula. *Queensland Naturalist*, 28 (1-4): 22-30

Covacevich, J.A. and Couper, P.J. 1991 Atlas of Queensland's frogs, reptiles, birds and mammals. Part 1.2 The reptile records. [Pp. 45-140]. In: Ingram, G.J. and Raven, R.J.

(Editors): Atlas of Queensland's frogs, reptiles, birds and mammals. Queensland Museum, Brisbane

Covacevich, J.A. and McDonald, K.R. 1991 Frogs and reptiles of tropical and subtropical eastern Australian rainforests: Distribution patterns and conservation. [Pp. 281-309]. In: Werren, G. and Kershaw, P. (Editors): The Rainforest Legacy. Australian National Rainforests Study. Volume 2. Flora and Fauna of Rainforests. Special Australian Heritage Publication Series, 7 (2) [Australian Government Publishing Service, Canberra]

Covacevich, J.A. and McDonald, K.R. 1991 Reptiles. [Pp. 69-82]. In: Nix, H.A. and Switzer, M.A. (1991): Rainforest Animals. Atlas of vertebrates endemic to Australia's wet tropics. Kowari, No 1

Covacevich, J.A., Davie, P. and Pearn, J. (Editors) 1987 Toxic Plants and Animals. A Guide for Australia. Queensland Museum, Brisbane

Coventry, A.J. 1970 Reptile and amphibian Type Specimens housed in the National Museum of Victoria. *Memoirs of the National Museum of Victoria*, 31: 115-124

Coventry, A.J. and Robertson, P. 1991 The Snakes of Victoria. Victorian Department of Conservation and Environment, Melbourne

Creed, J.M. 1887 Fear as a factor in producing many of the alarming symptoms following the bite of Australian snakes. [Pp. 36-46]. In: Transactions of the Intercolonial Medical Congress of Australasia, first session held in Adelaide, South Australia, August-September 1887. Vardon and Pritchard, Adelaide

Creed, J.M. 1887 Fear as a factor in producing many of the alarming symptoms following the bite of Australian snakes. Webb, Vardon and Pritchard, Adelaide [undated, but published ca. 1887; reprinted from Transactions of the Intercolonial Medical Congress of Australasia [Adelaide], for 1887; this is in effect, a revised Edition of Creed, J.M. (1884), of 12 pp.]

Crommelin, C.E. 1896 Case of snake-bite (Brown Snake) - Fatal. *Australasian Medical Gazette*, 15: 231-232

Davies, W. (Editor) 1983 Wildlife of the Brisbane Area. Jacaranda Press, Milton

Davis, R. and Mirtschin, P.J. 1985 Snake bites: A guide to treatment. *Patient Management*, 1985 (September): 91-99

De Boer, L.E.M. and Wissen, B. van 1978 The reptiles, amphibians and fish of Australia and New Guinea. [Pp. 75-80]. In: DeBoer, L.E.M. and Wissen, B. van (1978): A historical zoogeographic account of Australia, New Guinea and New Zealand. *Notogaea* [Rotterdam], Zoological Essays, 1: 1-96

De Vis, C.W. 1884 Descriptions of a new snake with a synopsis of the genus *Hoplocephalus*. *Proceedings of the Royal Society of Queensland*, 1: 138-140

De Vis, C.W. 1911 Description of snakes, apparently new. *Annals of the Queensland Museum*, 10: 22-25

- Doery, H.M. and Pearson, J.E. 1961 Haemolysins in venoms of Australian snakes. Observations on the haemolysins of the venoms of some Australian snakes and the separation of Phospholipase-A from the venom of *Pseudechis porphyriacus*. *Biochemical Journal*, 78: 820-827
- Douglas, A.M. and Ride, W.D.L. 1962 The results of an expedition to Bernier and Dorre Islands, Shark Bay, Western Australia. *Reptiles*. Western Australia Fisheries Department, Fauna Bulletin [Perth], 2: 113-119
- Dumeril, A.M.C., Bibron, G. and Dumeril, A.H.A. 1854 *Erpetologie Generale ou Histoire Naturelle Complete des Reptiles*. Roret, Paris [Volume 7, part 2; pp. 781-1536]
- Edgar, B. and Stephens, S. 1993 Commonwealth legislation relevant to reptiles and amphibians. [Pp. 39-42]. In: Lunney, D. and Ayers, D. (Editors): *Herpetology in Australia: A diverse discipline*. Royal Zoological Society of New South Wales, Sydney [an unnumbered 'Transactions of the Royal Zoological Society of New South Wales']
- [Edwards, H.V.] and Wild-Life Photography. How to obtain valuable snapshots of birds, beasts, reptiles, etc, in their native haunts. With 16 illustrations (13 from original photographs), an account of Australian Snipe, and the story of some interesting snapshots. Gordon and Gotch Ltd., Sydney [Pp. 1-32]
- Ehmann, H.F.W. 1976 The reptiles of the Mt Lofty Ranges, South Australia. Part 1. *Herpetofauna*, 8 (1): 2-5
- Ehmann, H.F.W. 1976 The reptiles of the Mt Lofty Ranges, South Australia. Part 2. *Herpetofauna*, 8 (2): 5-13
- Ehmann, H.F.W. 1992 *Encyclopaedia of Australian Wildlife*. Reptiles. Australian Museum, Sydney [Pp. 1-495]
- Fairley, N.H. 1929 The dentition and biting mechanism of Australian snakes. *Medical Journal of Australia*, (1929) 1 (10): 313-327
- Fairley, N.H. 1929 The dentition and biting mechanism of Australian snakes. In: Fairley, N.H., Kellaway, C.H. and Splatt, B. (1929): *Symposium on Snake Bite*. *Medical Journal of Australia*, March, 1929: 19-33 [Symposium Proceedings]
- Fairley, N.H. and Splatt, B. 1929 Venom yields in Australian poisonous snakes. *Medical Journal of Australia*, (1929) 1 (11): 336-348
- Fairley, N.H. and Splatt, B. 1929 Venom yields in Australian poisonous snakes. In: Fairley, N.H., Kellaway, C.H. and Splatt, B. (1929): *Symposium on Snake Bite*. *Medical Journal of Australia*, March, 1929 : 34-36 [Symposium Proceedings]
- Fischer, J.G. 1856 *Neue Schlangen des Hamburgischen Naturhistorischen Museum*. *Abh.Geb. Natur. Hamb.*, 3: 79-116
- Fitzinger, L. 1860 Ausbeute der osterreichischen Naturforscher an Saugethiere und Reptilien wahrend der Weltumsegelung Sr. Majestat Fregatte Novara. *Sber. Akad. Wiss. Wien*, 42: 384-415

- Fleay, D.H. 1943 The Brown Snake (*Demansia textilis*) - Dangerous fellow. *Victorian Naturalist*, 59 (9): 147-152
- Frauca, H. 1982 What Animal is That? A Guide to Australian Amphibians, Insects, Mammals, Reptiles and Spiders. Reed, Sydney [Note that another Edition appeared in 1985 also]
- Freeman, M. and Kellaway, C.H. 1934 The venom yields of common Australian poisonous snakes in captivity. *Medical Journal of Australia*, (1934) 21 (12): 373-377
- Frith, C.B. and Frith, D.W. 1987 Australian Tropical Reptiles and Frogs. Tropical Australian Graphics, Paluma [Qld]
- Fry, D.B. 1914 On a collection of reptiles and batrachians from Western Australia. *Records of the Western Australian Museum and Art Gallery*, 1: 174-210
- Furneaux, R.W. 1967 Successful recovery in a dog from a Common Brown Snake (*Demansia textilis*) bite. *Australian Veterinary Journal*, 43: 147-148
- Gillam, M.W. 1979 The genus *Pseudonaja* (Serpentes: Elapidae) in the Northern Territory. Territory Parks and Wildlife Commission [Alice Springs], Research Bulletin, No 1: 1-28
- Glasby, C.J., Ross, G.J.B. and Beesley, P.L. (Editors) 1993 Fauna of Australia. Volume 2A. Amphibia and Reptilia. Australian Government Publishing Service, Canberra [Pp. 1-439]
- Glauert, L. 1950 A Handbook of the Snakes of Western Australia. Western Australian Naturalist's Club, Perth [printed by Paterson Brokensha, Perth; Pp. 1-48 + (2)]
- Glauert, L. 1957 A Handbook of the Snakes of Western Australia. Western Australian Naturalist's Club, Perth [printed by Paterson Brokensha, Perth; 2nd Edition; Pp. 1-62; Reprinted 1960]
- Glauert, L. 1967 A Handbook of the Snakes of Western Australia. Western Australian Naturalists' Club, Perth [3rd Edition; published posthumously]
- Golay, P. 1985 Checklist and Keys to the Terrestrial Proteroglyphs of the World. Elapsoidea, Geneva
- Golay, P. et al (Editors) 1993 Endoglyphs and other Major Venomous Snakes of the World. A Checklist. Elapsoidea, Geneva
- Gow, G.F. 1976 Snakes of Australia. Angus and Robertson, Sydney [Pp. 1-88]
- Gow, G.F. 1976 Snakes of the Darwin area. Northern Territory Museum, Darwin
- Gow, G.F. 1982 Australia's Dangerous Snakes. Angus and Robertson, London [Pp. 1- 80 + plates]
- Gow, G.F. 1983 Snakes of Australia. Angus and Robertson, London [2nd Edition ('Completely Revised Edition')]

Gow, G.F. 1989 Graeme Gow's Complete Guide to Australian Snakes. Angus and Robertson, Sydney

Gray, J.E. 1841 A catalogue of the species of Reptiles and Amphibia hitherto described as inhabiting Australia, with a description of some new species from Western Australia, and some remarks on their geographical distribution. [Pp. 422-449]. In: Grey, G. (1841): Journals of Two Expeditions of Discovery in North-west and Western Australia, during the years 1837, 38 and 39, under the Authority of Her Majesty's Government describing many newly discovered, important, and fertile districts, with Observations on the Moral and Physical Conditions of the Aboriginal Inhabitants &c, &c. T. and W. Boone, London [volume 2; Appendix E]

Gray, J.E. 1842 Catalogue of the species of reptiles and amphibia hitherto described as inhabiting Australia. Tasmanian Journal of Natural Science, Agriculture and Statistics, 1: 385-387

Gray, J.E. 1845 The reptiles of Australia. [Pp. 1-8 + plates]. In: Richardson, J. and Gray, J.E. (Editors): The Zoology of the Voyage of H.M.S. Erebus and Terror, under the Command of Capt. J. C. Ross during 1839-1845. Volume 2. Reeve Brothers, London

Green, D. 1973 The reptiles of the outer north-western suburbs of Sydney. Herpetofauna, 6 (2): 2-5

Green, K. and Osborne, W. 1994 Wildlife of the Australian Snow Country. Reed Books, Sydney

Greer, A.E. 1997 The Biology and Evolution of Australian Snakes. Surrey Beatty and Sons, Sydney

Greer, A.E. and Shine, R. 2000 Relationship between mass and length in Australian Elapid snakes. Memoirs of the Queensland Museum, 45 (2): 375-380

Griffiths, K. 1987 Reptiles of the Sydney Region. Three Sisters Productions, Winmalee

Griffiths, K. 1996 Australia's Reptiles and Frogs. Topmill Press, Sydney

Griffiths, K. 1997 Frogs and Reptiles of the Sydney Region. University of New South Wales Press, Kensington [Pp. 1-128]

Gunther, A. 1858 Catalogue of Colubrine Snakes in the Collection of the British Museum. British Museum, London

Gunther, A. 1872 Seventh account of new species of snakes in the collection of the British Museum. Annals and Magazine of Natural History, (4) 9: 13-37

Hawdon, G. and Winkel, K.D. 1997 Could this be snake bite? Australian Family Physician, 26: 1386-1391

Herrmann, R.P., Davey, M.G. and Skidmore, P.H. 1972 The coagulation defect after envenomation by the bite of the Dugite (*Demansia nuchalis affinis*), a Western Australian brown snake. Medical Journal of Australia, (1972) 2: 183-186

- Hobson, H.R. 1961 Brown Snake and Stumpy-tail Lizard. *Victorian Naturalist*, 77: 259
- Hoser, R.T. 1980 Further records of aggregations of various species of Australian snakes. *Herpetofauna*, 12 (1): 16-22
- Hoser, R.T. 1981 Reptiles of the Pilabara [sic] region, Western Australia. *Journal of the Northern Ohio Association of Herpetologists (NOAH)*, 7 (1): 12-32 [authorship erroneously attributed to 'Robert' Hoser]
- Hoser, R.T. 1986 Reptile populations of the Gunnedah and Boggabri districts, New South Wales, Australia. *Journal of the Northern Ohio Association of Herpetologists (NOAH)*, 12 (1): 19-29
- Hoser, R.T. 1988 Conservation of Australian snakes, other reptiles and frogs. *Litteratura Serpentina*, 8 (1): 12-40
- Hoser, R.T. 1989 Australian Reptiles and Frogs. Pierson and Co., Mosman [Pp. 1-238]
- Hoser, R.T. 1990 Melbourne's snakes, Part 1. *Litteratura Serpentina*, 10 (2): 82-92
- Hoser, R.T. 1990 Melbourne's snakes, Part 2. *Litteratura Serpentina*, 10 (3): 122-145
- Hoser, R.T. 1990 Pairing behaviour in Australian snakes. *Herpetile*, 15 (3): 84-92
- Hoser, R.T. 1990 Some experiences when photographing reptiles - the stories behind the photos. *Herpetile*, 15 (4): 128-136a
- Hoser, R.T. 1991 Sydney's snakes, Part 1. *Litteratura Serpentina*, 11 (3): 56-67
- Hoser, R.T. 1991 Sydney's snakes, Part 3. *Litteratura Serpentina*, 11 (5): 101-113
- Houston, T.F. 1973 Reptiles of South Australia. A brief synopsis. *South Australian Yearbook*, 1973 [11 pp.]
- Hunt, R.A. 1947 A key to the identification of Australian snakes (other than Blind Snakes, Rock Snakes and Pythons, File Snakes and Sea Snakes). *Victorian Naturalist*, 64 (8): 153-167
- Hutchinson, M. 1990 The generic classification of the Australian terrestrial Elapid snakes. *Memoirs of the Queensland Museum*, 29: 397-405
- Jacobson, K. 1973 Reptiles of the Tamworth area. *Herpetofauna*, 6 (1): 20-22
- Jan, G. 1859 Plan d'une Iconographie descriptive des Ophidiens, et description sommaire de nouvelles especes de Serpents. *Rev. Mag. Zool.*, (2) 11-12: 122-130
- Jan, G. 1859 Prodrome d'une Iconographie descriptive des Ophidiens, et Description sommaire de nouvelles especes de Serpents venimeux. Bouchard-Huzard, Paris
- Jan, G. 1863 Elenco Sistematico Degli Ofidi Descritti e Disegnati per l'Iconografia Generale. A. Lombardi, Milan [Pp. 1-143]

- Jan, G. and Sordelli, F. 1873 *Iconographie Generale des Ophidiens*. Atlas 1860-1881. Authors, Milan
- Jenkins, R.W.G. and Bartell, R.J. 1980 *A Field Guide to Reptiles of the Australian High Country*. Inkata Press, Melbourne [Pp. 1-187]
- Johnson, C.R. 1975 Defensive display behaviour in some Australian and Papuan-New Guinean Pygopodid lizards, Boid, Colubrid and Elapid snakes. *Zoological Journal of the Linnean Society*, 56: (4): 265-282
- Johnston, G.R. and Ellins, P. 1979 The reptiles of the Sir Joseph Banks Islands, South Australia. *Herpetofauna*, 10 (2): 9-12
- Johnston, T.H. and Mawson, P.M. 1948 Some new records of nematodes from Australian snakes. *Records of the South Australian Museum*, 9: 101-106
- Jones, H.I. 1978 *Abbreviata* (Nematoda: Physalopteroidea) from Western Australian snakes. *Australian Journal of Zoology*, 26 (4): 789-808
- Jones, H.I. 1980 Observations on nematodes from west and central Australian snakes. *Australian Journal of Zoology*, 28 (3): 423-433
- Jones, P. 1990 A record of predation on *Nephrurus levis* by *Pseudonaja nuchalis*. *Herpetofauna*, 20 (1): 15-16
- Kay, P., Garlepp, M. and Dawkins, R.L. 1979 Evidence for human anti-venom antibodies and detection of anti-acetylcholine receptor antibodies using Dugite (*Pseudonaja affinis*) venom. In: Chubb, I.W. and Geffen, L.B. (Editors): *Neurotoxins and Clinical Advances*. Centre for Neuroscience, Flinders Medical Centre, Adelaide
- Keast, J.A. 1959 The reptiles of Australia. [Pp. 115-135]. In: Keast, A., Crocker, R.L. and Christian, C.S. (Editors): *Biogeography and Ecology in Australia*. W. Junk, The Hague [Monographiae Biologicae, 8]
- Keast, J.A. 1962 Vertebrate speciation in Australia: Some comparisons between Birds, Marsupials and Reptiles. [Pp. 380-407]. In: Leeper, G.W. (Editor): *The Evolution of Living Organisms*. A symposium to mark the centenary of Darwin's 'Origin of Species' and of the Royal Society of Victoria held in Melbourne, December 1959. Melbourne University Press, Melbourne
- Kellaway, C.H. 1931 Observations on the certainly lethal dose of the venom of the Common Brown Snake (*Demansia textilis*) for the common laboratory animals. *Medical Journal of Australia*, (1931) 2: 747-751
- Kellaway, C.H. 1931 The immunity of Australian snakes to their own venoms. *Medical Journal of Australia*, (1931) 2: 35-52
- Kellaway, C.H. 1932 Venomous land snakes in Australia. *Bulletin of the Antivenin Institute of America*, 5: 53-58

- Kellaway, C.H. 1934 The venoms of some of the small and rare Australian venomous snakes. *Medical Journal of Australia*, (1934) 2: 74-78
- Kellaway, C.H. 1934 Alleged antidotes in the treatment of the bites of Australian snakes. *Melbourne Hospital Clinical Reports*, 5: 4-12
- Kellaway, C.H. 1938 The symptomatology and treatment of the bites of Australian snakes. *Medical Journal of Australia*, (1938) 2: 585-589
- Kellaway, C.H. 1942 The symptomatology and treatment of the bites of Australian snakes. *Medical Journal of Australia*, (1942) 2: 171-174
- Kellaway, C.H. and Eades, T. 1929 Field notes on the common Australian venomous snakes. *Medical Journal of Australia*, (1929) 2: 249-257
- Kellaway, C.H. and Williams, F.E. 1931 The serological and blood relationships of some common Australian snakes. *Australian Journal of Experimental Biology and Medical Science*, 8: 123-132
- Kendrick, P.G. and Rolfe, J.K. 1991 The reptiles and amphibians of Kimberley rainforests. [Pp. 347-359]. In: McKenzie, N.L., Johnston, R.B. and Kendrick, P.G. (Editors): *Kimberley Rainforests of Australia*. Surrey Beatty and Sons, Chipping Norton [Sydney]
- Kinghorn, J.R. 1920 Studies in Australian reptiles. 1. *Records of the Australian Museum*, 13 (3): 110-117
- Kinghorn, J.R. 1924 Reptiles and batrachians from south and south-west Australia. *Records of the Australian Museum*, 14: 163-183
- Kinghorn, J.R. 1929 *The Snakes of Australia*. Angus and Robertson, Sydney [Pp. 1-198 + 122 plates]
- Kinghorn, J.R. 1945 Australian poisonous snakes. *Australian Museum Magazine*, 8 (10): 325-330
- Kinghorn, J.R. 1949 Birds and reptiles. In: Poignant, A. (Editor): *Bush Animals of Australia*. Shepherd Press, Sydney
- Kinghorn, J.R. 1951 Reptiles and amphibians. [Pp. 774-788]. In: *The Australian Junior Encyclopaedia*. Georgian House, Melbourne
- Kinghorn, J.R. 1956 *The Snakes of Australia*. Angus and Robertson, Sydney [2nd Edition; first published 1929; note: see also reprinting of 1964]
- Kinghorn, J.R. 1968 Reptiles. [Pp. 1-125]. In: *Animals of the World: Australia*. Hamlyn Publishing, Sydney
- Klemmer, K. 1963 Liste der Rezenten Giftschlangen: Elapidae, Hydrophidae, Viperidae und Crotalidae. [Pp. 255-464]. In: *Die Giftschlangen der Erde*. N.G. Elwert Univ., Marburg

- Krefft, J.L.G. 1862 Notes upon Australian snakes, and their geographical distribution. Proceedings of the Zoological Society of London, 1862: 224-226
- Krefft, J.L.G. 1869 The Snakes of Australia; An illustrated and descriptive catalogue of all the known species. Government Printer, Sydney [Pp. i-xxv, 1-100]
- Lee, C.-Y. (Editor) 1979 Venoms. Handbook of Experimental Pharmacology. 52: 922-955
- Lester, J.W. and Bellairs, A. d'A. 1954 Australian reptiles. Zoo Life, 9: 50-54
- Lillywhite, H.B. 1980 Behavioral thermoregulation in Australian Elapid snakes. Copeia, 1980: 452-458
- Longman, H.A. 1915 Reptiles from Queensland and the Northern Territory. Memoirs of the Queensland Museum, 3: 30-34
- Longman, H.A. 1926 Snakes. [Pp. 470-475]. In: The Australian Encyclopaedia. Volume 2. Angus and Robertson, Sydney
- Longmore, R. (Editor) 1986 Atlas of Elapid Snakes of Australia. Bureau of Flora and Fauna, Canberra [Australian Flora and Fauna Series, Number 7; Pp. i-v, 1-115; see also 2nd Edition]
- Lonnberg, E. and Andersson, G.L. 1913 Results of Dr E. Mjobergs' Swedish Scientific Expeditions to Australia 1910-1913. III. Reptiles. Kunglia Svenska Vetenskapsakademiens Handlingar, 52 (3): 1-17
- Lonnberg, E. and Andersson, G.L. 1915 Results of Dr E. Mjobergs' Swedish Scientific Expeditions to Australia 1910-1913. VII. Reptiles collected in northern Queensland. Kunglia Svenska Vetenskapsakademiens Handlingar, 52 (7): 1-9
- Loveridge, A. 1934 Australian reptiles in the Museum of Comparative Zoology, Cambridge, Massachusetts. Bulletin of the Museum of Comparative Zoology [Harvard], 77 (6): 243-383
- Loveridge, A. 1938 On some reptiles and amphibians from the central region of Australia. Transactions of the Royal Society of South Australia, 62 (2): 183-191
- Low, T. 1978 The reptiles of Magnetic Island, north Queensland. Herpetofauna, 9 (2): 10-14
- Lowry, E. 1964 The venomous snakes of Australia. Australian Journal of Pharmacy, 45: 527-531
- Lucas, A.H.S. and LeSouef, W.H.D. 1909 The Animals of Australia. Mammals, Reptiles and Amphibians. Whitcombe and Tombs, Melbourne [Pp. i-xi, 1-327]
- Lunney, D. and Barker, J. 1986 Survey of reptiles and amphibians of the coastal forests near Bega, NSW. Australian Zoologist, 22 (3): 1-9
- Maaten, F. and Stevens, A. 1982 Some observations on captive breeding of Western Brown Snakes. Wildlife Australia, 19 (2): 42-43

- Mack, G. and Gunn, S.B. 1953 'De Vis' Types of Australian Snakes. *Memoirs of the Queensland Museum*, 13 (1): 58-70
- Mackerras, M.J. 1961 The haematozoa of Australian reptiles. *Australian Journal of Zoology*, 9: 61-122
- Markwell, K. and Knight, R. 1986 An inventory of the herpetofauna of the south-eastern section of Myall Lakes National Park, NSW. Hunter Wetlands Trust [Newcastle], Scientific Paper No 1
- Macleay, W.J. 1884 Census of Australian snakes with descriptions of two new species. *Proceedings of the Linnean Society of New South Wales*, 9: 548-568
- Macleay, W.J. 1885 On a new snake from the Barrier Ranges. *Proceedings of the Linnean Society of New South Wales*, 10: 61-62
- Martin, C.J. 1896 An explanation of the marked difference in the effects produced by subcutaneous and intravenous injection of the venom of Australian snakes. *Journal and Proceedings of the Royal Society of New South Wales*, 30: 150-157
- Martin, C.J. 1897 The curative value of Calmette's anti-venomous serum in the treatment of inoculations with the poisons of Australian snakes. *Intercolonial Medical Journal of Australasia*, 2 (6): 527-536
- Martin, C.J. 1898 The curative value of Calmette's anti-venomous serum in the treatment of inoculations with the poisons of Australian snakes. (A rejoinder to M. le Dr. Calmette). *Intercolonial Medical Journal of Australasia*, 3: 197-204
- McCann, I.R. 1983 *Little Desert Wildlife: An illustrated checklist*. Little Desert Tours, Pty Ltd, Nhill
- McCoy, F. 1867 On the recent zoology and palaeontology of Victoria. *Annals and Magazine of Natural History*, (3) 20: 175-202
- [McCoy, F.] 1870 Our Colonial Snakes. No 3. Brown, Black and Tiger Snakes. *Australasian* [Melbourne], (ns) January 1: 8 [Authored By 'Microzoon']
- McCoy, F. 1879 *Diemenia superciliosa* (Fischer). The Common Brown Snake. *Prodromus of the Zoology of Victoria; or, figures and descriptions of the living species of all Classes of the Victorian indigenous animals. Decade III: Plate 23, Figure 1 + 2 pp.*
- McCoy, F. 1879 The Shield-fronted Brown Snake. *Prodromus of the Zoology of Victoria; or, figures and descriptions of the living species of all Classes of the Victorian indigenous animals. Decade III: 13-14*
- McCoy, F. 1879 The Two-hooded Furina Snake. *Prodromus of the Zoology of Victoria; or, figures and descriptions of the living species of all Classes of the Victorian indigenous animals. Decade IV: 13-15, Plate 4, fig 2]*

- McDermid, E.M., Board, P.G. and Agar, N.S. 1977 Studies on the blood of Australian Elapid snakes. III. Electrophoretic analysis of serum proteins and red cell enzymes. *Comparative Biochemistry and Physiology*, (B) 56: 361-365
- McDowell, S.B. 1967 *Aspidomorphus*, a genus of New Guinea snakes of the family Elapidae, with notes on related genera. *Journal of Zoology* [London], 151: 497-543
- McDowell, S.B. 1985 The terrestrial Australian Elapids: General summary. [Pp. 261-264]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- McPhee, D.R. 1959 *Some Common Snakes and Lizards of Australia*. Jacaranda, Brisbane [Pp. 1-125]
- McPhee, D.R. 1963 *Some Common Snakes and Lizards of Australia*. Jacaranda, Brisbane [Second Edition; Pp. 1-125]
- McPhee, D.R. 1979 *The Observer's Book of Snakes and Lizards of Australia*. Methuen, Sydney [Pp. 1-157]
- Mengden, G.A. 1984 The taxonomy of Australian Elapid snakes: A review. *Records of the Australian Museum*, 35: 195-222 [Note: This work although dated 1983, was not published until 1984]
- Mengden, G.A. 1985 A chromosomal and electrophoretic analysis of the genus *Pseudonaja*. [Pp. 193-208]. In: Grigg, G., Shine, R. et al (Editors): *The Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Mengden, G.A. 1985 Australian Elapid phylogeny: A summary of the chromosomal and electrophoretic data. [Pp. 185-192]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Meyer, G. 1977 Reptiles of 'Timbertop', Mt Bulla, Victoria. *Victorian Herpetological Society Newsletter*, 5 (September): 2-3
- Milhinch, A.L. 1967 Observations on a Gwardar (*Demansia nuchalis*). *Western Australian Naturalist*, 10 (7): 169-170
- Miller, B. and Schwaner, T.D. 1983 The Speckled Brown Snake *Pseudonaja guttata* Parker: An addition to the fauna of South Australia. *Transactions of the Royal Society of South Australia*, 106 (2): 79-80
- Mincham, H. 1970 *Reptiles of Australia and New Zealand*. Rigby, Adelaide [Pp. 1-63]
- Minton, S.A. and Minton, M.R. 1981 Toxicity of some Australian snake venoms for potential prey species of reptiles and amphibians. *Toxicon*, 19 (6): 749-755

- Mirtschin, P.J. and Davis, R. 1982 Dangerous snakes of Australia - An illustrated guide to Australia's most venomous snakes. Rigby, Adelaide [Pp. 1-207]
- Mirtschin, P.J. and Davis, R. 1983 Dangerous Snakes of Australia. An illustrated guide to Australia's most venomous snakes. Rigby, Adelaide ['Revised' Edition]
- Mirtschin, P.J. and Davis, R. 1991 Dangerous Snakes of Australia: An Illustrated Guide to Australia's most Venomous Snakes. Ure Smith, Sydney [2nd Edition]
- Mirtschin, P.J. and Davis, R. 1992 Snakes of Australia: Dangerous and Harmless. Hill of Content Publishers, Melbourne [Pp. 1-216]
- Mirtschin, P.J., Crowe, G.R. and Davis, R. 1990 Dangerous snakes of Australia. In: Gopalakrishnakone, P. and Chou, L.M. (Editors): Snakes of Medical Importance (Asia-Pacific Region). Venom and Toxin Research Froup, National University of Singapore and the International Society of Toxinology (Asia-Pacific Region)
- Mitchell, F.J. 1951 The South Australian reptile fauna. Part 1. Ophidia. Records of the South Australian Museum, 9: 545-557
- Mitchell, F.J. 1958 Adaptive convergence in Australian reptiles. Australian Museum Magazine, 12 (10): 314-317
- Morgan, F.G. 1929 Preparation of curative sera against venom of certain Australian snakes. [Pp. 466-469]. In: Archdall, M. (Editor): Australasian Medical Congress Transactions of the Third Session Sydney. Government Printer, Sydney [see also Abstract in Medical Journal of Australia, 2: 537 (1929)]
- Munday, B.L., Hartley, W.J., Harrigan, K.E., Presidente, P.J.A. and Obendorf, D.L. 1979 Sarcocystis and related organisms in Australian wildlife: II. Survey findings in Birds, Reptiles, Amphibians and Fish. Journal of Wildlife Diseases, 15 (1): 57-74
- Nix, H.A. 1986 A biogeographic analysis of Australian Elapid snakes. [Pp. 4-15]. In: Longmore, R. (Editor) (1986): Atlas of Elapid Snakes of Australia. Bureau of Flora and Fauna, Canberra [Australian Flora and Fauna Series Number 7]
- O'Shea, M. 1996 A Guide to the Snakes of Papua New Guinea. Independent Publishing, Port Moresby [Pp. i-xii, 1-238]
- Owner, F. 1970 Reptiles of Australia and New Zealand. Angus and Robertson, Sydney [Pp. 1-30]
- Parker, H.W. 1926 New reptiles and a new frog from Queensland. Annals and Magazine of Natural History, (9) 17: 665-670
- Peters, U.W. 1971 Der tod in Sydney's Garten. Die Braunschlange, Pseudonaja t. textilis ist ein kulturfolger. Aquarien-Magazin [Stuttgart], 5: 36
- Rankin, P.R. 1973 Lizard mimicking a snake - Juvenile Tiliqua casuarinae (Dumeril and Bibron). Herpetofauna, 5 (4): 13-14

- Rawlinson, P.A. 1965 Snakes of the Melbourne area. *Victorian Naturalist*, 81: 245-254
- Rawlinson, P.A. 1966 Reptiles of the Victorian mallee. *Proceedings of the Royal Society of Victoria*, 79: 605-619
- Rawlinson, P.A. 1969 The reptiles of East Gippsland. *Proceedings of the Royal Society of Victoria*, 82: 113-128
- Rawlinson, P.A. 1971 The reptiles of West Gippsland. *Proceedings of the Royal Society of Victoria*, 84: 37-51
- Reilly, V.M. 1960 Three poisonous brown snakes. *North Queensland Naturalist*, 28 (125): 9-10
- Riley, J. and Spratt, D.M. 1987 Further observations on Pentastomids (Arthropoda) parasitic in Australian reptiles and mammals. *Records of the South Australian Museum*, 21 (2): 139-147
- Riley, J., Spratt, D.M. and Presidente, P.J.A. 1985 Pentastomids (Arthropoda) parasitic in Australian reptiles and mammals. *Australian Journal of Zoology*, 33: 39-53
- Roberts, J. and Mirtschin, P.J. 1991 An uncommon prey record for the Common Brown Snake *Pseudonaja textilis*. *Herpetofauna*, 21 (1): 36 [Published in December, 1991 - not May, 1991 as printed on cover]
- Robinson, A.C. 1980 Notes on the mammals and reptiles of Pearson, Dorothee and Greenly Islands, South Australia. *Transactions of the Royal Society of South Australia*, 104 (5): 93-99
- Rose, A.B. 1974 Gut contents of some amphibians and reptiles. *Herpetofauna*, 7 (1): 4-8
- Saint-Girons, H. and Bradshaw, S.D. 1981 Preliminary observations on behavioural thermoregulation in an Elapid snake, the Dugite, *Pseudonaja affinis* Gunther. *Journal of the Royal Society of Western Australia*, 64: 13-16
- Schapel, G.J., Utley, D. and Wilson, G.C. 1971 Envenomation by the Australian Common Brown Snake - *Pseudonaja (Demansia) textilis textilis*. *Medical Journal of Australia*, (1971) 1: 142-144
- Schmida, G. 1986 *The Cold Blooded Australians. A unique photographic study of Australia's reptiles, amphibians and freshwater fish.* Doubleday, Sydney
- Schwaneer, T.D., Baverstock, P.R., Dessauer, H.C. and Mengden, G.A. 1985 Immunological evidence for the phylogenetic relationships of Australian Elapid snakes. [Pp. 177-184]. In: Grigg, G., and Shine, R. et.al. (Editors): *Biology of Australasian Frogs and Reptiles.* Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Schwaneer, T.D., Miller, B. and Tyler, M.J. 1985 Reptiles and amphibians. In: Tyler, M.J. (Editor): *Natural History of Eyre Peninsula.* Royal Society of South Australia, Adelaide

- Seddon, H.D. 1974 Snakebite in a cat by a Five-Ringed Snake (*Demansia modesta*). *Australian Veterinary Journal*, 50: 326-327
- Shine, R. 1975 Ecological Studies on Australian Elapid Snakes (Ophidia; Elapidae). PhD Thesis, University of New England, Armidale [Pp. 1-254]
- Shine, R. 1977 Reproduction in Australian Elapid snakes. I. Testicular cycles and mating seasons. *Australian Journal of Zoology*, 25: 647-653
- Shine, R. 1977 Reproduction in Australian Elapid snakes. II. Female reproductive cycles. *Australian Journal of Zoology*, 25: 655-666
- Shine, R. 1977 Habitats, diet, and sympatry in snakes: A study from Australia. *Canadian Journal of Zoology*, 55: 1118-1128
- Shine, R. 1978 Sexual size dimorphism and male combat in snakes. *Oecologia*, 33: 269-278
- Shine, R. 1979 Activity patterns in Australian Elapid snakes (Squamata, Serpentes, Elapidae). *Herpetologica*, 35 (1): 1-11
- Shine, R. 1981 Niche relationships in Australian snakes. [p.143]. In: Banks, C.B. and Martin, A.A. (Editors): *Proceedings of the Melbourne Herpetological Symposium*. Zoological Board of Victoria, Melbourne [Abstract only]
- Shine, R. 1985 Ecological evidence on the phylogeny of Australian Elapid snakes. [Pp. 255-260]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Shine, R. 1985 The reproductive biology of Australian reptiles: A search for general patterns. [Pp. 297-303]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Shine, R. 1989 Constraints, allometry, and adaptation: Food habits and reproductive biology of Australian Brownsnakes (*Pseudonaja*: Elapidae). *Herpetologica*, 45 (2): 195-207
- Shine, R. 1991 *Australian Snakes: A Natural History*. Reed Books, Sydney [Pp. 1-223]
- Shine, R. and Covacevich, J. 1983 Ecology of highly venomous snakes: The Australian genus *Oxyuranus* (Elapidae). *Journal of Herpetology*, 17 (1): 60-69
- Simpson, K.N.G. 1973 Amphibians, reptiles and mammals of the Murray River region between Mildura and Renmark, Australia. *Memoirs of the National Museum of Victoria*, 34: 275-279
- Slater, K.R. 1968 *A Guide to the Dangerous Snakes of Papua*. Bloink, Port Moresby
- Smith, L.A. 1976 The reptiles of Barrow Island. *Western Australian Naturalist*, 13 (6): 125-136

- Smith, L.A. and Johnstone, R.E. 1981 Amphibians and reptiles of Mitchell Plateau and adjacent coast and lowlands, Kimberley, Western Australia. [Pp. 215-227] In: Biological Survey of Mitchell Plateau and Admiralty Gulf, Kimberley, Western Australia. Western Australian Museum, Perth
- Smith, M.J. 1975 The vertebrae of four Australian Elapid snakes (Squamata: Elapidae). Transactions of the Royal Society of South Australia, 99 (2): 71-84
- Sonter, C.R. and Debus, S.J.S. 1985 The Brown Falcon, *Falco berigora* as a predator of snakes. Australian Bird Watcher, 11: 92-93
- Stackhouse, J. 1970 Australia's Venomous Wildlife. Paul Hamlyn, Sydney [Pp. 1-144]
- Steindachner, F. 1867 Reptilien. In: Reise der osterreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859. Zoologie, 1 (3): 1-98
- Stevens, N.C. and Bailey, A. (Editors) 1980 Contemporary Cape York Peninsula. Royal Society of Queensland, Brisbane [Pp. 1-100]
- Stopford, F. 1995 Profile of an Eastern Brown Snake *Pseudonaja textilis*. Hawkesbury Herpetologist, 7: 18-20
- Storr, G.M. 1964 Some aspects of the geography of Australian reptiles. Senckenbergiana Biologia, 45 (3-5): 577-589
- Storr, G.M. 1979 Dangerous Snakes of Western Australia. Western Australian Museum, Perth [3rd Edition; Pp. 1-24]
- Storr, G.M. 1985 Phylogenetic relationships of Australian Elapid snakes: External morphology with an emphasis on species in Western Australia. [Pp. 221-222]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): Biology of Australasian Frogs and Reptiles. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Storr, G.M. 1989 A new *Pseudonaja* (Serpentes: Elapidae) from Western Australia. Records of the Western Australian Museum, 14 (3): 421-423
- Storr, G.M. and Smith, L.A. 1975 Amphibians and Reptiles of the Prince Regent River Reserve, north-western Australia. [Pp. 85-88]. In: Miles, J.M. and Burbidge, A.A. (Editors): A Biological Survey of the Prince Regent River Reserve, north-west Kimberley, Western Australia. Western Australian Wildlife Research Bulletin, No 3: 1-116
- Storr, G.M. and Smith, L.A. 1977 Amphibians and Reptiles. [Pp. 97-101]. In: Kabay, E.D. and Burbidge, A.A. (Editors): A Biological Survey of the Drysdale River National Park, North Kimberley, Western Australia in August 1975. Western Australian Wildlife Research Bulletin, No 6
- Storr, G.M. and Smith, L.A. 1981 Amphibians and Reptiles. [Pp. 54-59]. In: Mackenzie, N.L. (Editor): Wildlife of the Edgar Ranges area, south-west Kimberley, Western Australia. Western Australian Department of Fisheries and Wildlife, Wildlife Research Bulletin, No 10: 1-71

- Storr, G.M., Smith, L.A. and Johnstone, R.E. 1986 Snakes of Western Australia. Western Australian Museum, Perth [Pp. 1-187]
- Sutherland, S.K. 1979 Snakebites and antivenoms. *Australian Prescriber*, 3: 84-87
- Sutherland, S.K. 1981 *Venomous Creatures of Australia. A field guide with notes on first aid.* Oxford University Press, Melbourne [Pp. i-xi, 1-131]
- Sutherland, S.K. 1983 *Australian Animal Toxins. The creatures, their toxins and care of the poisoned patient.* Oxford University Press, Melbourne
- Sutherland, S.K. 1992 Deaths from snake bite in Australia, 1981-1991. *Medical Journal of Australia*, 157: 740-745
- Sutherland, S.K. and King, K. 1991 *Management of snakebite in Australia.* Royal Flying Doctor Service of Australia, Monograph Series No 1
- Swan, G. 1990 *A Field Guide to the Snakes and Lizards of New South Wales.* Three Sisters Publications, Winmalee [Pp. 1-224]
- Swan, G. 1998 *Green Guide. Snakes and Other Reptiles of Australia.* New Holland, Sydney [Pp. 1-96]
- Tamiya, N. 1985 A comparison of amino acid sequences of neurotoxins and of phospholipases of some Australian Elapid snakes with those of other Proteroglyphous snakes. [Pp. 209-219]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles.* Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Tate, R. 1881 South Australian snakes and lizards, Notes on. *Transactions of the Royal Society of South Australia*, 4: 139
- Thomas, W.D. 1982 From Goannas to Taipans. The reptiles of Australia. *Zooview*, 16 (2): 5, 16
- Thomson, D.F. and Hosmer, W. 1963 A preliminary account of the herpetology of the Great Sandy Desert of central Western Australia. *Reptiles and amphibians of the Bindibu Expedition. Proceedings of the Royal Society of Victoria*, 77 (1): 217-237
- Trethewie, E.R. 1971 The pharmacology and toxicology of the venoms of the snakes of Australia and Oceania. [Pp. 79-101]. In: Bucherl, W. and Buckley, E.E. (Editors): *Venomous Animals and Their Venoms. Volume 2.* Academic Press, New York
- Underhill, D. 1987 *Australia's Dangerous Creatures. Reader's Digest*, Sydney
- Underwood, G. 1967 *A Contribution to the Classification of Snakes*, British Museum, London
- Waite, E.R. 1898 *A Popular Account of Australian Snakes with a Complete List of the Species and an Introduction to their Habits and Organisation.* Thomas Shine, Sydney [Pp. 1-71 + 16 Plates]

- Waite, E.R. 1916 The 'deadly' snakes of South Australia. *Children's Hour for Reading and Recreation*, 28 (320): 114-118
- Waite, E.R. 1925 Field notes on some Australian reptiles and batrachians. *Records of the South Australian Museum*, 3: 17-32
- Waite, E.R. 1927 The fauna of Kangaroo Island, South Australia. No 3. The reptiles and amphibians. *Transactions of the Royal Society of South Australia*, 51: 326-329
- Waite, E.R. 1929 The Reptiles and Amphibians of South Australia. Government Printer, Adelaide [Published posthumously; Handbooks of the Flora and Fauna of South Australia, Edited by Hale, H.M. and issued by the British Science Guild, South Australian Branch, Adelaide; Pp. 1-270 + 192 figures; see also facsimile reprint of 1993 by SSAR]
- Wallach, V. 1985 A cladistic analysis of the terrestrial Australian Elapidae. [Pp. 223-253]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Watson, W.G. 1908 Australian snakes. *Journal of the American Medical Association*, 50: 293
- Webb, G.A. and Rose, A.B. 1986 The food of some Australian snakes. *Herpetofauna*, 16 (1): 21-27 [not 1984 as printed on cover]
- Weigel, J. 1988 Care of Australian Reptiles in Captivity. Reptile Keepers Association, Gosford [Pp. 1-143]
- Weigel, J. 1990 The Australian Reptile Park's Guide to the Snakes of South-East Australia. Australian Reptile Park, Gosford [Pp. 1-103]
- Wells, R.W. 1980 Eggs and young of *Pseudonaja textilis textilis*. *Herpetofauna*, 12: 30-32
- Wells, R.W. and Turner, J.R. 1988 Checklist of the reptiles, frogs and Jewel Beetles of the Agnes Banks area, New South Wales, Australia. *Australian Herpetologist*, No 515: 1-9
- Wells, R.W. and Wellington, C.R. 1984 Synopsis of the Class Reptilia in Australia. *Australian Journal of Herpetology*, 1 (3-4): 63-129
- Wells, R.W. and Wellington, C.R. 1985 A Classification of the Amphibia and Reptilia in Australia. *Australian Journal of Herpetology (Supplementary Series)*, No 1: 1-61
- Wells, R.W. and Wellington, C.R. 1988 A checklist of the amphibians and reptiles known from the lower Blue Mountains region, Sydney Basin, New South Wales, Australia. *Australian Herpetologist*, No 508: 1-7
- Wells, R.W. and Wellington, C.R. 1988 The amphibians and reptiles of the Blue Mountains region, Sydney Basin, New South Wales, Australia. *Australian Herpetologist*, No 504: 1-12
- Wells, R.W. and Wellington, C.R. 1988 Preliminary checklist of the amphibians and reptiles of the Kanangra Plateau, Sydney Basin, New South Wales, Australia. *Australian Herpetologist*, No 507: 1-3

- Wells, R.W. and Wellington, C.R. 1988 Amphibians and reptiles of the upper Cox's River area, Sydney Basin, New South Wales, Australia, with comments on Greer and Cogger's recent reclassification of the genus *Anomalopus* (sensu lato). *Australian Herpetologist*, No 505: 1-15
- Wells, R.W. and Wellington, C.R. 1989 A Checklist of the Amphibians and Reptiles known from the Cumberland Plain Region, Sydney Basin, New South Wales, Australia. *Australian Herpetologist*, 506: 1-34
- West, A.B. 1960 Brown snake's meal. *Victorian Naturalist*, 76 (11): 288-289
- White, J. 1981 Ophidian envenomation: A South Australian perspective. *Records of the Adelaide Children's Hospital*, 2: 311-421
- White, J., Pounder, D., Pearn, J.H. and Morrison, J.J. 1985 A perspective on the problems of snakebite in Australia. [Pp. 511-514]. In: Grigg, G., Shine, R. and Ehmann, H. (Editors): *Biology of Australasian Frogs and Reptiles*. Surrey Beatty and Sons, Sydney [Note that the publication date is erroneously given as August, 1985 - actually published in November 1985]
- Williams, F.E., Freeman, M. and Kennedy, E. 1934 The bacterial flora of the mouths of Australian venomous snakes in captivity. *Medical Journal of Australia*, (1934) 2: 190-193
- Wilson, S.K. and Knowles, D.G. 1988 *Australia's Reptiles - A photographic reference to the terrestrial reptiles of Australia*. Collins, Melbourne [Pp. 1-447]
- Wilson, S.K. and Swan, M. 1972 A study of the reptiles of the Hattah Lakes district. *Junior Naturalist*, 8 (3): 5-11
- Worrell, E. 1952 Snakes of Australia. *Outdoors and Fishing* [Sydney], 9 (4): 236-237
- Worrell, E. 1952 Our dangerous snakes. *Snakes of Australia*. Part 3. *Outdoors and Fishing* [Sydney], 10 (1): 38-39
- Worrell, E. 1952 *Dangerous Snakes of Australia*. Angus and Robertson, Sydney [64 pp.]
- Worrell, E. 1953 *Dangerous Snakes of Australia*. Angus and Robertson, Sydney [2nd Edition, 64 pp.]
- Worrell, E. 1956 Notes on skull characters of some Australian snakes. *Australian Zoologist*, 12 (3): 205-210
- Worrell, E. 1957 *Dangerous Snakes of Australia*. Angus and Robertson, Sydney [3rd Edition, 68 pp.]
- Worrell, E. 1961 A new insular Brown Snake. *Proceedings of the Royal Zoological Society of New South Wales for 1958-59*: 56-58
- Worrell, E. 1961 *Dangerous Snakes of Australia and New Guinea*. Angus and Robertson, Sydney [Usually regarded as the 4th Edition of Worrell (1952), however it should be regarded as the first edition of a distinct publication in its own right, as the title, text and illustrations were substantially altered; 66 pp]

Worrell, E. 1963 Dangerous Snakes of Australia and New Guinea. Angus and Robertson, Sydney [2nd Edition of Worrell (1961), note also that it was reprinted in 1966; 66pp.]

Worrell, E. 1963 Reptiles of Australia: Crocodiles - Turtles - Tortoises - Lizards - Snakes. Describing all Australian species, their appearance, their haunts, their habits, with over 330 illustrations, many in full colour. Angus and Robertson, Sydney [Pp. i-xv + 1-207]

Worrell, E. 1966 Australian Snakes, Crocodiles, Tortoises, Turtles, Lizards. Angus and Robertson, Sydney [Pp. 1-64]

Worrell, E. 1969 Dangerous Snakes of Australia and New Guinea. Angus and Robertson, Sydney [3rd Edition of Worrell (1963), 65 pp.]

Worrell, E. 1970 Reptiles of Australia: Crocodiles - Turtles - Tortoises - Lizards - Snakes. Describing their appearance, their habits, with over 330 illustrations, many in full colour. Angus and Robertson, Sydney [2nd Edition, without synonymic checklist; Pp. i-xv + 1-169]

Worrell, E. 1969 Dangerous Snakes of Australia and New Guinea. Angus and Robertson, Sydney

[Note: This list of references is only a selection of the available publications that deal with the species mentioned in this paper. If readers would like a larger listing they are invited to write to the author of this paper]

EDITORIAL NOTICE

The Australian Biodiversity Record is published at no cost for permanent scientific record in accordance with the International Commission of Zoological Nomenclature Rules (Code 4th Edition). Copies may be obtained by sending a written request to be placed on a mailing list. The address for submission of manuscripts, subscriptions and all other enquiries relating to this Journal is:

Australian Biodiversity Record
"Shiralee", Major West Road,
Cowra, NSW Australia 2794

Note: Although the articles published by the Australian Biodiversity Record always remain the Intellectual Property of the authors concerned, it is an expressed condition of publication that articles may be freely copied by any person and distributed worldwide by any means whatsoever for the sole purpose of research or criticism. All articles so copied from the Australian Biodiversity Record must include full bibliographic data that identifies the author name/s, the journal name and number, and date of publication. Therefore, the Australian Biodiversity Record recommends that any article copied be a full and complete copy to ensure that due credit is given to authors and to also allow their information to be considered in accurate context.

Pseudonaja genus. accepted (GBIF) listed (PBDB). 4. Cope E. D. On the homologies of some of the cranial bones of the Reptilia, and on the systematic arrangement of the class, Proceedings of the American Association for the Advancement of Science 19, 194-247.

Cope E. D. Check-list of North American Batrachia and Reptilia; with a systematic list of the higher groups, and an essay on geographical distribution. Based on the specimens contained in the U. S. National Museum, Bulletin of the United States National Museum 1, 1-109.

Zittel K. A. v. , Handbuch der Palaeontologie. I. Abteilung Paleozoologie. III. Band. Vertebrata (Pisces, Amphibia, Reptil... Pseudonaja. Quite the same Wikipedia. Just better. Pseudonaja is a genus of venomous elapid snakes native to Australia. Species of this genus are known commonly as brown snakes and are considered to be some of the most dangerous snakes in the world; even young snakes are capable of delivering a fatal envenomation to a human. Despite its common name, the king brown snake (Pseudechis australis) is not a brown snake, but a member of the genus Pseudechis, commonly known as black snakes.

Contents. 1 Species.