

Preliminary Report of a Controlled Trial of MTH-68/BTM Virus Vaccine Treatment in Acute B and C Hepatitis: A Phase II Study

LASZLO K. CSATARY¹, LÁSZLÓ TELEGDY², PÉTER GERGELY³, BELA BODEY⁴
and TIBOR BAKÁCS⁵

¹United Cancer Research Institute, Alexandria, VA, U.S.A.;

²St. Laszlo Hospital, Budapest, Hungary;

³Semmelweis University, Budapest Hungary;

⁴University of Southern California, Los Angeles, CA, U.S.A.;

⁵National Institute of Oncology, Budapest, Hungary

Abstract. Eighty four patients with viral hepatitis attributed to infection with hepatitis B virus (HBV) (n=43) or hepatitis C virus (HCV) (n=41) were included in this study employing the MTH-68/BTM vaccine, an attenuated variant of Bursal Disease Virus. Twenty of the 43 patients in the HBV group, and 22 of the 41 HCV patients were treated with MTH-68/BTM. The remaining patients received conventional therapy. Significantly more patients progressed into active chronic hepatitis on conventional therapy (13% of HBV and 26% of HCV cases respectively) than in the vaccine treated groups (0% and 9%). Relapses occurred less frequently in the vaccine treated groups (5% of HBV and 32% of HCV) than in the control groups (9% and 79%), while remissions within one month of treatment were observed more often in the vaccine treated groups (both 50% respectively) than in the control groups (26% of HBV and 21% of HCV patients respectively). The duration of hepatitis was also considerably shortened by MTH-68/BTM treatment in both HBV (from 7.5±3.7 to 5.9±3.0 weeks) and HCV patient groups (from 8.9±7.4 to 5.3±4.4 weeks). The data presented suggest that attenuated, non-pathogenic viruses may be of significant benefit for patients with viral hepatitis B and C infections.

Viral hepatitis is one of the most common infectious disease of humans. It is an inflammation of the liver which is caused by different viruses (hepatitis A, B, C, D, E and G respectively). Based on human studies and animal models of HBV pathogenesis, there is considerable evidence that viral hepatitis is initiated by a specific antiviral cellular immune

response, but most of the damage to the liver is caused by a cascade of non-specific effector systems induced by the infection (1). In the majority of cases this disease appears in an acute form, and most of the patients recover completely. Five to 10 percent of acute HBV infections but 60-80% of HCV infections tend to progress into chronic active hepatitis in which the liver cells are gradually destroyed (2). The role of chronic B and C hepatitis in the etiology of liver cirrhosis and various malignancies, including primary hepatocellular carcinoma has been established (3-7). Chronic infection can occur without clinical symptoms of acute hepatitis, since HBV often infects individuals, particularly infants and children, who remain asymptomatic carriers of virus sometimes for decades. Following such an eclipse phase, chronic hepatitis and primary HCC may develop. As the estimated number of HBV carriers is more than 350 millions, it is not surprising that HCC is ranking within the ten most common cancers of the world (8).

The prevalence of hepatitis infection can be significantly decreased by the vaccination of children. Universal vaccination has therefore been adopted in more than 70 countries (9). Although the full efficiency of such vaccination programs on the occurrence of hepato-carcinoma will be seen only some 40 years later, it already has been demonstrated in Taiwan that the incidence of hepatocellular carcinoma in children has significantly declined (9).

Until the full effect of such vaccination programs will be felt, however, successful treatment of acute B and C hepatitis remains an important medical problem. Despite significant efforts in the development of treatment strategies for viral hepatitis, as yet there is little that can be done to prevent progression into chronic infection. Treatment of chronic viral hepatitis with interferon- α and other antiviral drugs is not only very expensive, but unsuccessful in many cases. Therefore, it is clear that special efforts must be made to prevent chronic hepatitis (10).

MTH-68/BTM is an attenuated variant of Bursal Disease

Correspondence to: Tibor Bakács, 1122, National Institute of Oncology, Budapest, Ráth György u. 7-9. Hungary; tel.: 361-1554411/ext.1165; Fax: 361-1562402; E-mail: bakacs@oncol.hu

Key Words: Acute viral hepatitis, hepatitis B virus (HBV), hepatitis C virus (HCV), hepatocellular carcinoma (HCC), bursal disease virus (BDV), virus vaccine therapy, phase II study.

Table I. Comparison of MTH-68/BTM virus vaccine and conventional treatment of viral hepatitis B and C.

	Hepatitis B		Hepatitis C	
	Vaccine	Control	Vaccine	Control
Progression into Chronic Active Hepatitis	0/20 (0%)	3/23* (13%)	2/22 (9%)	5/19* (26%)
Occurrence of Relapses	1/20 (5%)	2/23 (9%)	7/22 (32%)	15/19*** (79%)
Occurrence of Late Remissions (over 6 months)	0/20 (0%)	4/23** (17%)	3/22 (14%)	8/19** (42%)
Remission within 1 month	10/20 (50%)	6/23 (26%)	11/22 (50%)	4/19 (21%)
Duration of Hepatitis (weeks±SD)	5.9 ± 3.0	7.5 ± 3.7	5.3 ± 4.4	8.9 ± 7.4

Significance (p value, chi-square, Yates' correction): *p < 0.05; **p < 0.02; ***p < 0.01

Virus (BDV), which is a double-stranded RNA virus and a member of the Birnaviridae family (11). As this virus is apathogenic for humans, it has no public health significance (12). An attenuated avian bursa virus was successfully used in the treatment of human hepatitis A viral infection in marmoset monkeys (13). More importantly, when the MTH-68/BTM vaccine has been administered in terminal cancer patients occasionally rapid resolution of viral hepatitis was observed (14). To evaluate the effect of MTH-68/BTM treatment in acute hepatitis B and C, a Phase II trial was conducted data of which are presented here.

Patients and Methods

84 patients of both sexes (ages 14-70 years) with either a diagnosis of acute B (43 patients) or acute C (41 patients) viral hepatitis were included in this study. They were hospitalized because of jaundice, other clinical signs of acute hepatitis (fever, severe malaise, loss of appetite), and a 10-100 fold elevation of alanine-aminotransferase (ALAT) level. The diagnosis of hepatitis B viral (HBV) infection was verified by the presence of HBsAg, HBeAg, and anti-HBeIgM antibody. Acute hepatitis C viral (HCV) infection was determined by the exclusion of A and B virus, EBV, and CMV infection and by the appearance of anti-HCV antibody. Exclusion criteria from this study were the presence of viral markers other than hepatitis B and C, respectively, HIV positivity, suspicion or evidence of alcohol- or drug-induced hepatitis, clinical or histological signs of chronic hepatitis as well as other chronic liver diseases, fulminant hepatitis, malignancies, underlying systemic disease, immunosuppressive treatment within 6 months, pregnancy, and lack of compliance. Patients were randomly allocated into two groups upon admittance. One group was treated with MTH-68/BTM vaccine, while the other received conventional treatment.

The trial protocol was approved by the National Health Scientific Council and the Ethical Committee of St. László Hospital. All patients gave their informed consent. The studies were performed according to the Helsinki Declaration of Human Rights. The patients received MTH-68/BTM vaccine, a live attenuated BDV (Phylaxia, Hungary) intranasally once every day (4000 U/day) for 1 week, then 3 times a week for two weeks, and finally once a month for 6 months. Clinical and laboratory investigations were performed before treatment, weekly for one month and once every month for 6 months consisting of physical examination, detection of viral markers, liver function tests (bilirubin, ALAT, ASAT,

alkaline phosphatase, γ -glutamyl transferase) and others (ESR, blood count, urinalysis). Liver biopsy was optional (in case of suspicion of chronic hepatitis). Side effects were registered. Criteria of *remission*: normalization of serum bilirubin and ALAT levels, disappearance of HBsAg, no relapse within 6 months; *relapse*: elevation of ALAT (more than two times of upper normal limit) with or without return of jaundice after cessation of therapy within 6 months; *subacute course*, late *remission*: prolonged elevation (more than two times of normal upper limit) of ALAT after the 6th month of the disease, no viral seroconversion; *chronic course*: more than two-fold elevation of ALAT level after one year, histological signs of chronic hepatitis at percutaneous liver biopsy, HBsAg and/or anti-HBe IgM or HCV-PCR positivity, respectively. Patients with progression into chronic hepatitis and viraemia proven by PCR were treated with interferon- α after one year. Hepatitis serology was tested by commercial enzyme immunoassay kits (Organon and Ortho Microelisa).

Statistical analysis was performed by the Yates' correction of chi-square test.

Results and Discussion

In the present study, an attenuated form of BDV, (MTH-68/BTM virus vaccine) was used for the treatment of viral hepatitis B and C. Consistently with the literature, more patients progressed into chronic hepatitis in the HCV than HBV group. However, there was a significant difference between the vaccine treated and control groups, as only 9% of HCV and none of the HBV patients progressed into chronic disease, whereas 13% and 26% of the controls did (Table I). Prior to complete recovery 9% of HBV and 79% of HCV control patients, but only 5% and 32% of vaccine treated patients relapsed. Late remissions (requiring more than 6 months) were recorded significantly more frequently with conventional treatment in both HBV and HCV hepatitis groups (17% and 42% respectively in controls but only 0% and 14% with vaccine treatment). MTH-68/BTM treatment also shortened the duration of the first icteric phase (by 20% in HBV and 40% in HCV groups). Furthermore more remission within one month of treatment was registered in the virus treated groups (both 50% respectively) than with

conventional treatment (26% and 21%). No side effects of virus vaccine treatment were noted.

It has been shown earlier that concurrent viral infections may significantly influence the outcome of viral disease in humans (15,16,17). Recent studies using transgenic mouse models of the hepatitis B virus infection have revealed how an unrelated viral infection of the liver can be 'curative' to hepatitis B (18). It was demonstrated that hepatocellular HBV gene expression and replication can be abolished by noncytopathic, cytokine-dependent pathway (principally by TNF- α and IFN- α/β produced by intrahepatic macrophages) which were activated by an unrelated lymphocytic choriomeningitis virus (LCMV) infection of the liver (19). These results imply, that coinfection or superinfection of the HBV-infected liver by other pathogens may facilitate HBV clearance if they induce the local production of cytokines such as IFN γ or TNF α (20). In accordance with this some case report presented evidence that chronic HBV infection resolved during an intermittent infection of the liver by hepatitis A or C (20,21). Conceptually similar events may be operative in other viral infections as well. Indeed, control of viral replication by IFN α/β , IFN γ and TNF α has been suggested for LCMV, vaccinia, measles, herpes simplex, influenza, Semliki Forest and Theiler's viruses respectively (22-31). It is tempting to speculate that MTH-68/BTM vaccine treatment may resolve the viral hepatitis by a similar mechanism. In conclusion, the MTH-68/BTM virus vaccine therapy reported in this study seems to be safe and efficient in treating acute HBV and HCV mediated hepatitis, thus it may be of significant importance in the prevention of the chronic disease. To reach unequivocal conclusions and possibly elucidate the underlying mechanism of action of the vaccine treatment, we believe, that further investigation is warranted.

References

- Chisari FV, Ferrari C: Hepatitis B virus immunopathogenesis. *Annu Rev Immunol* 13: 29-60, 1995.
- Tsai JF, Chang WY, Jeng JE, Ho MS, Lin ZY, Tsai JH: Hepatitis B and C virus infection as risk factors for liver cirrhosis and hepatocellular carcinoma: A case-control study. *Liver* 14: 98-102, 1994.
- Hoofnagle JH: Hepatitis B: *In: Bokus Gastroenterology* (Haubrich WS, Schaffner F, and Berk JE, eds). Saunders, Philadelphia, pp. 2026-2081, 1995.
- Tsai JF, Chang WY, Jeng JE, Ho MS, Wang LY, Hsieh MY, Chen SC, Chuang WL, Lin ZY, Tsai JH: Hepatitis C virus infection as a risk factor for non-alcoholic liver cirrhosis in Taiwan. *J Med Virol* 41: 296-300, 1993.
- Tsai JF, Chang WY, Jeng JE, Ho MS, Lin Zy, Tsai JH: Frequency of raised α -foetoprotein level among Chinese patients with hepatocellular carcinoma related to hepatitis B and C. *Br J Cancer* 69: 1157-1159, 1994.
- Tsai JF, Jeng JE, Ho MS, Chang WY, Lin ZY, Tsai JH: Hepatitis B and C virus infection as risk factors for hepatocellular carcinoma in Chinese: A case-control study. *Int J Cancer* 56: 619-621, 1994.
- Tsai JF, Chang WY, Jeng JE, Ho MS, Lin ZY, Tsai JH: Effects of hepatitis C and B viruses infection on the development of hepatocellular carcinoma. *J Med Virol* 44: 92-95, 1994.
- Blumberg BS: Hepatitis B virus, the vaccine, and the control of primary cancer of the liver. *Proc Natl Acad Sci USA* 94: 7121-7125, 1997.
- Chang MH, Chen CJ *et al*: For the Taiwan Childhood Hepatoma Study Group: Universal Hepatitis B vaccination in Taiwan and the incidence of hepatocellular carcinoma in children. *The N Eng J Med* 336: 1855-1858, 1997.
- Viral Hepatitis Management. Standards for the future. Abstract Book, Cannes, Macclesfield, UK: Adelphi Communications, Ltd, 1992.
- Kibens FS, Dhillon AS, Russell RG: Biochemistry of infectious bursal disease virus. *J Gen Virol* 69: 1757-1775, 1988.
- Peterson KA, Sadasiv EC, Chang PW, Yates VJ: Detection of antibody to avian viruses in human populations. *Epidemiol Infect* 104: 519-525, 1990.
- Csatary LK, Kasza L, Massey RJ: Interference between human hepatitis A virus and an attenuated apathogenic avian virus. *Acta Microbiol Hung* 31: 153-158, 1984.
- Csatary LK, Gergely P: Virus vaccines for the treatment of cancer. *Orv Hetil* 131: 2585-2588, 1990.
- Csatary LK: Viruses in the treatment of cancer. *Lancet* II: 285, 1971.
- Csatary LK, Romvary JJ, Toth B, Tauber LN: *In vivo* study of interferences between herpes and influenza viruses. *J Med* 13: 1-7, 1982.
- Csatary LK, Romvary JJ, Kasza L, Schaff S, Massey RJ: *In vivo* interference between pathogenic and non-pathogenic viruses. *J Med* 16: 563-573, 1985.
- Guidotti LG, Chisari FV: To kill or to cure: options in host defense against viral infection. *Curr Opin Immunol* 8: 478-483, 1996.
- Guidotti LG, Borrow BD, Hobbs MV, Matzke B, Gresser I, Oldstone MBA, Chisari FV: Viral cross talk: intracellular inactivation of the hepatitis B virus during an unrelated viral infection of the liver. *Proc Natl Acad Sci USA* 93: 4589-4594, 1996.
- Davis GL, Hoofnagle JH, Waggoner JG: Acute type A hepatitis during chronic hepatitis B virus infection: association of depressed hepatitis B virus replication with appearance of endogenous alpha interferon. *J Med Virol* 14: 141-147, 1984.
- Sheen IS, Liaw YF, Lin DY, Chu CM: Role of hepatitis C and delta viruses in the termination of chronic hepatitis B surface antigen carrier state: a multivariate analysis in a longitudinal follow-up study. *J Infect Dis* 170: 1358-1361, 1994.
- Muller U, Steinhoff U, Reis LFL, Hemmi S, Pavlovic J, Zinkernagel RM, Aguet M: Functional role of type I and II interferons in antiviral defense. *Science* 264: 1918-1921, 1994.
- Sambhi SK, Kohonen-Corish MRJ, Ramshaw IA: Local production of tumor necrosis factor encoded by recombinant vaccinia virus is effective in controlling viral replication *in vivo*. *Proc Natl Acad Sci USA* 88: 4025-4029, 1991.
- Kohonen-Corish MRJ, King NJC, Woodhams CE, Ramshaw IA: Immunodeficient mice recover from infection with vaccinia virus expressing interferon- γ . *Eur J Immunol* 20: 157-161, 1990.
- Van Den Broek MF, Muller U, Huang S, Aguet M, Zinkernagel RM: Antiviral defense in mice lacking both alpha/beta and gamma interferon receptors. *J Virol* 69: 4792-4796, 1995.
- Finke D, Brinckmann UG, Ter Meulen V, Liebert UG: Gamma interferon is a major mediator of antiviral defense in experimental measles virus-induced encephalitis. *J Virol* 69: 5469-5474, 1995.

- 27 Rossol-Voth R, Rossol S, Schutt KH, Corridori S, De Cian W, Falke D: *In vivo* protective effect of tumor necrosis factor α against experimental infection with herpes simplex virus type I. *J Gen Virol* 72: 143-147, 1991.
- 28 Bouley DM, Kanangat S, Wire W, Rouse BT: Characterization of herpes simplex virus type-1 infection and herpetic stromal keratitis development in IFN- γ knockout mice. *J Immunol* 155: 3964-3971, 1995.
- 29 Topham DJ, Tripp RA, Sarawar SR, Sangster MY, Doherty PC: Immune CD4⁺ T cells promote the clearance of influenza virus from major histocompatibility complex class II^{-/-} respiratory epithelium. *J Virol* 70: 1288-1291, 1996.
- 30 Hwang SY, Hertzog PJ, Holland KA, Sumarsono SH, Tymms MJ, Hamilton JA, Whitty G, Bertoncello I, Kola I: A null mutation in the gene encoding a type I interferon receptor component eliminates anti-proliferative responses to interferons α and β and alters macrophage responses. *Proc Natl Acad Sci USA* 92: 11284-11288, 1995.
- 31 Fiette L, Aubert C, Muller U, Huang S, Aguet M, Brahic M, Bureau JF: Theiler's virus infection of 129Sv mice that lack the interferon α/β or interferon γ receptors. *J Exp Med* 181: 2069-2076, 1995.

Received November 17, 1997

Accepted December 22, 1997

Preliminary results of an NIH-sponsored clinical trial (ACTT; NCT04280705) published in NEJM showed remdesivir accelerated recovery from advanced COVID-19.[7]. Preliminary Results from ACTT trial . Remdesivir. A Multicenter, Adaptive, Randomized Blinded Controlled Trial of the Safety and Efficacy of Investigational Therapeutics for the Treatment of COVID-19 in Hospitalized Adults. ClinicalTrials.gov NCT04280705. [<https://clinicaltrials.gov/ct2/show/NCT04280705>]. ARDS Definition Task Force, Ranieri VM, Rubenfeld GD, et al.