

Stability constants of binary and ternary complexes of some transition metal ions

A.B. PATIL

Department of Chemistry, P.N. College, Pusad - 445 216 (India)

(Received: April 12, 2008; Accepted: July 05, 2008)

ABSTRACT

The stability constants of Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) metal ions with pyridoxine hydrochloride (PHC) and ethambutol hydrochloride (EHC) has been studied by pH-metric technique at 0.1 M (KNO_3) ionic strength at 302 ± 0.5 K in aqueous medium. The data obtained were used to evaluate the values of proton-ligand and metal-ligand stability constants using Irving-Rossotti titration technique. Mixed-ligand complex studies of these metal ions using aspartic acid (ASP) and glutamic acid (GLU) as primary ligands and PHC and EHC as secondary ligands also have been carried out pH-metrically at the same conditions.

The Irving-Williams order of stability constants was observed in case of binary as well as ternary complexes.

Key words: Ternary complexes, transition metal ions, pyridoxine hydrochloride, ethambutol hydrochloride, aspartic acid and glutamic acid

INTRODUCTION

Recently there has been considerable interest in the study of binary, ternary and quaternary complexes by pH-metric method¹⁻³. The ligand pyridoxine hydrochloride (PHC) is a vitamin. It plays a part in protein metabolism, the synthesis of fat from protein, haemopoiesis and the nutrition of skin⁴⁻⁶. The ligand ethambutol hydrochloride (EHC) is used in the treatment of mycobacterial infection, particularly tuberculosis⁴⁻⁶. Ternary complexes of Ni (II) and Cu (II) with nicotinic acid as primary ligand and imidazoles, benzimidazole, histamine and L-histidine as secondary ligands have been studied by Nair and Neekantan⁷. Patil and Mhaske⁸ have studied the stability constants of Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) with nitrilotriacetic acid and Iminodiacetic acid as primary ligands and pyridoxine hydrochloride and ethambutol hydrochloride as secondary ligands potentiometrically. Nigam and coworkers⁹ have studied the ternary complexes of Mn (II), Co (II), Ni

(II), Cu (II) and Zn (II) using ASP as primary ligand and thymine as secondary ligand potentiometrically. Ternary complexes of Cu (II) using ASP and GLU as primary ligands have been reported potentiometrically by Pandeya and Patel¹⁰. In this paper the stability constants of binary and ternary complexes of Mn (II), Co (II), Ni (II), Cu (II) and Zn (II) ions with ASP and GLU as primary ligands and PHC as secondary ligand at 302 ± 0.5 K and at fixed ionic strength, $\mu = 0.1$ M KNO_3 using modified form of Irving-Rossotti pH-metric technique¹¹ in aqueous medium have been studied.

EXPERIMENTAL

The ligand PHC was obtained from LOBA Chemie and the EHC was of Cadila Pharmaceutical Ltd. The ligands were used as such. Carbonate free sodium hydroxide solution was prepared by standard method¹². All other solutions were prepared in doubly distilled water.

The pH-metric measurements were carried out by using Elico digital pH-meter model L-120 with combined glass-calomel electrode with an accuracy of ± 0.01 of pH unit at 302 ± 0.5 K. the pH-meter was standardized against 0.05 M potassium hydrogen phthalate solution in acid medium and 0.01 M borax solution in alkaline medium.

For determination of proton-ligand stability constant of the secondary ligand and the metal-ligand stability constants of binary and ternary complexes, the following set of solutions were prepared and titrated against standard alkali solution.

Binary Systems

- 9.6×10^{-3} M HNO_3
- 9.6×10^{-3} M HNO_3 + 5.0×10^{-3} M secondary ligand.
- 9.6×10^{-3} M HNO_3 + 5.0×10^{-3} M secondary ligand + 1.0×10^{-3} M metal ion.

Ternary Systems

- 9.6×10^{-3} M HNO_3
- 9.6×10^{-3} M HNO_3 + 1.0×10^{-3} M secondary ligand
- 9.6×10^{-3} M HNO_3 + 1.0×10^{-3} M primary

- ligand + 1.0×10^{-3} M metal ion
- 9.6×10^{-3} M HNO_3 + 1.0×10^{-3} M primary ligand + 1.0×10^{-3} M metal ion + 1.0×10^{-3} M secondary ligand.

The ionic strength was maintained constant (0.1 M) by adding required volume of 1M KNO_3 . The ratio of metal (M): secondary ligand (L) was maintained at 1: 5 in each of the binary systems and the ratio of metal (M): primary ligand (A): secondary ligand (L) was maintained at 1: 1: 1 in each of the ternary system.

RESULTS AND DISCUSSION

Proton-Ligand Stability Constants

The plots of volume of alkali (NaOH) against pH-meter readings were used to evaluate the proton-ligand stability constants of PHC and EHC. The deviation between free acid titration curve and secondary ligand titration curve was used to evaluate the formation functions, \bar{n}_A . The proton-ligand formation curves were then obtained by plotting the values of \bar{n}_A versus pH-meter readings. From the graphs, the values of $\log K_1^H$ and $\log \beta^H$ were evaluated by half integral method (method A) and point wise calculation method (method B) and presented in Table 1.

Table 1: Proton-Ligand Stability Constants

Ligand	log		log K_1^H		log β^H	
	Method A	Method B	Method A	Method B	Method A	Method B
PHC	9.05	9.03	4.95	4.92	14.00	13.95
EHC	9.45	9.40	6.52	6.54	15.97	15.94

Metal-ligand stability constants of binary complexes

The metal-ligand stability constants of binary complexes were evaluated assuming that the formation of hydrolyzed products, polynuclear complexes, hydrogen and hydrogen bearing complexes were absent. An examination of titration curves indicates that complex formation has taken place in the solution on the following grounds:

1. The metal titration curves showed displacement with respect to the ligand

titration curves along the volume axis. This indicated the affinity of the ligand with metal ions which released protons and produced the volume difference ($V_3 - V_2$).

2. The colour change of the ligand in presence of metal ions appeared showing the formation of new species.
3. The hydrolysis of the metal ions was suppressed due to complex formation and the precipitation did not appear during the titration.

From the ligand and metal titration curves the value of $\bar{\eta}$ and from that the values of pL were obtained. The formation curves obtained were used to evaluate the metal-ligand stability constants by method (A) and (B) and presented in Table 2.

The variation of $\bar{\eta}$ was found to be 0 to 2 which indicated that the composition of complexes was 1:2 in solution.

The Irving-Williams order^{13, 14} of stability constants was followed.

Metal-ligand stability constants of ternary complexes

The metal-ligand stability constants of the ternary complexes were evaluated assuming that the formation of hydrolyzed products, polynuclear complexes, hydrogen and hydrogen bearing complexes were absent. An examination of titration

curves indicates that complex formation has taken place in the solution on the following grounds:

1. The ternary complex titration curves show the displacement with primary complex titration curves. The horizontal distance was measured between acid curve and the secondary ligand curve (V_2-V_1) and subtracted through the horizontal distance between ternary complex curves and primary complex titration curves (V_4-V_3) show the positive difference which proves the earlier released of protons in the formation of ternary complexes.
2. The hydrolysis of the metal ions was suppressed and precipitation did not result.

The values of $\bar{\eta}$ varies from 0 to 0.8, thus confirming the formation of 1: 1:1 mixed ligand complexes. The values of $\log K_{MAL}^{ASP}$ and $\log K_{MAL}^{GLU}$ have been evaluated from the formation curves ($\bar{\eta}$ vs. pL)

Table 2: Metal-Ligand Stability Constants of Binary Complexes


System	$\log K_1$		$\log K_2$		$\log \beta$	
	Method A	Method B	Method A	Method B	Method A	Method B
[Mn(II)-PHC]	2.62	2.62	-	-	-	-
[Co(II)-PHC]	2.92	2.96	2.75	2.75	5.67	5.71
[Ni(II)-PHC]	3.39	3.49	2.81	2.82	6.20	6.31
[Cu(II)-PHC]	5.08	5.06	4.62	4.63	9.70	9.69
[Zn(II)-PHC]	3.88	3.87	3.52	3.50	7.40	7.37
[Mn(II)-EHC]	4.18	4.17	3.28	3.30	4.46	4.37
[Co(II)-EHC]	5.65	5.67	3.68	3.65	9.33	9.32
[Ni(II)-EHC]	6.62	6.62	3.80	3.83	10.42	10.45
[Cu(II)-EHC]	6.92	6.93	4.74	4.71	10.66	10.64
[Zn(II)-EHC]	5.79	5.83	3.57	3.61	9.36	9.44

Table 3: Metal-Ligand Stability Constants of Ternary Complexes

System	Mn(II)		Co(II)		Ni(II)		Cu(II)		Zn(II)	
	A	B	A	B	A	B	A	B	A	B
[M (II)-ASP-PHC]	3.99	4.00	4.97	4.96	5.07	5.05	5.50	5.51	5.03	5.04
[M (II)-GLU-PHC]	3.96	3.99	4.40	4.42	5.51	5.48	5.81	5.87	4.73	4.75
[M (II)-ASP-EHC]	4.81	4.80	6.12	6.14	6.32	6.35	7.18	7.20	5.54	5.54
[M (II)-GLU-EHC]	4.58	4.60	6.15	6.16	6.48	6.47	6.92	6.96	6.23	6.24

At $\eta^- = 0.5$ in the formation curve, $pL = \log K$. The metal- ligand stability constant of PHC and EHC as secondary ligands and ASP and GLU as primary ligand are presented in Table 3.

The Irving-Williams order^{13, 14} of stability constants was observed in case of binary as well as ternary complexes which is



This sequence of stability of complexes with respect to metal ion is due to decreasing atomic radius and increasing the second ionization potential.

ACKNOWLEDGEMENTS

The authors are thankful to UGC and Principal, P. N, College, Pusad for providing necessary facilities.

REFERENCES

1. A. E. Martell and R. M. Smith, *Critical Constants-1: Amino acids*, New York (1974).
2. H. Sigel, *Metal Ions in Biological Systems-2*, Marcel-Dekker, Inc., New York (1973).
3. M. T. Beck, *Chemistry of Complex Equilibria*, Van Nostrand, New York, 174-190 (1970).
4. *The Pharmaceutical Codex*, 11th Edition, The Pharmaceutical Press, London (1979).
5. *Clarke's Isolation and Identification of Drugs*, 2nd Edition, The Pharmaceutical Press, London (1986).
6. *Remington's Pharmaceutical Science*, 17th Edition, Mack Publishing Co., Easton, Pennsylvania (1985).
7. M.S. Nair and M. A. Neelkantan, *Indian J. Chem.*, **38A**: 1307(1999).
8. A. B. Patil and T. H. Mhaske, *Asian J. Chem.*, **14**(1): 125-129 (2002).
9. N. B. Nigam, P. C. Sinha, M. Gupta and M. N. Srivastava, *Indian J. Chem.*, **22A**: 818 (1983).
10. K.B. Pandeya and R. N. Patel, *Indian J. Chem.*, **30A**: 193 (1991).
11. H. Irving and H. S. Rossotti, *J. Chem. Soc.*, **76**: 2904(1954).
12. A.I. Vogel, *A Text Book of Quantitative Analysis*, Longman, London, pp.241 (1961).
13. H. Irving and R.P.J. Williams, *Nature*, **162**: 746 (1948)
14. H. Irving and R.P.J. Williams, *J. Chem. Soc.*, **76**: 31

The stability constants of the 1:1 and 1:2 (whenever possible) complexes formed between La^{3+} , Ce^{3+} , Th^{4+} and the amino acid anions L-alaninate, L-phenylalaninate and L-histidinate were determined by potentiometric titration in aqueous solution ($25 \pm 1^\circ\text{C}$, $I = 0.1 \text{ M}$ KCl) and compared together with the constants previously determined. The various formation degree of the resulting M(L) and M(L)_2 were determined. In order to relate the formation degree of M(L) and M(L)_2 with the basicity of the amino acid anion (L^-), the acidity constants of the protonated amino acids, H_2L^+ , were also measured. The